



# Boards and Beyond: Infectious Disease

A Companion Book to the Boards and Beyond Website

Jason Ryan, MD, MPH

2018 Edition



# Table of Contents

Bacterial Cells	1	Protein Synthesis Inhibitors	91
Shapes and Stains	7	Other Antibiotics	97
Bacterial Culture	11	Fungal Pneumonias	102
Special Growth	15	Fungal Skin Infections	106
Virulence	18	Opportunistic Fungal Infections	109
Growth and Genetics	25	Antifungal Drugs	115
Bacterial Identification	29	Malaria	119
Staphylococci	33	Protozoa	123
Streptococci	37	Helminths	129
Other Gram Positives	42	Viral Structure	136
Gram Negative Rods	48	DNA Viruses	140
Other Gram Negatives	55	Herpes Viruses	145
Spirochetes	62	RNA Viruses	151
Zoonotic Infections	66	Viral Hepatitis	160
Mycobacteria	71	HIV	167
Penicillins	74	HIV Drugs	171
Beta Lactams	81	Antivirals	175
Sulfonamides	86		



# Bacteria

Jason Ryan, MD, MPH

## Types of Organisms

### Prokaryotes

- Very old form of life
- No membrane-bound organelles
- No nucleus
- Nuclear material free inside cell
- Bacteria are prokaryotes

### Eukaryotes

- More modern form of life
- Membrane-bound organelles
- Nucleus
- Plant and animal cells
- Protozoa
- Fungi

## Bacteria

- Single cell organisms
- Cell wall is key component
  - General support
  - Osmotic pressure
- Target for immune system
- Target for antibiotics
- Differentiates bacteria




Image courtesy Wikipedia/Public Domain

## Gram Stain

- Different for gram (+) and gram (-) bacteria





Image courtesy of Y tambe /Wikipedia

## Cell Walls

Lipoteichoic Acids



Gram Positive Bacteria

Cell membrane



Gram Negative Bacteria

## Peptidoglycan

- Major structural component of bacterial cell walls
- Polymer sheets of sugars and peptides
- Sheets cross-linked to other sheets

## Peptidoglycan

- Sugars:
  - *N*-acetylglucosamine (NAG)
  - *N*-acetylmuramic acid (NAM)
- Peptides:
  - Attached to NAM
  - Three to five amino acids
- Sugar/peptide backbone makes chains
- Chains cross-linked by peptide cross-bridges
- Site of action some antibiotics
  - Penicillin, cephalosporins

## Peptidoglycan




Image courtesy of Twooars / Wikipedia

## Peptidoglycan

- Gram positive bacteria
  - Up to 40 sheets
  - 50% or more of cell wall
- Gram negative bacteria
  - Very few sheets
  - 5-10% cell wall
- Thick layer in gram (+) bugs retains the gram stain
  - Makes them purple

## Unique Cell Walls

- Mycoplasma
  - No cell wall
  - Does not gram stain
  - Cell membrane has sterols for extra stability
- Mycobacteria
  - Cell wall has mycolic acid
  - Does not gram stain well
  - Special stains used (Ziehl-Neelsen)
- Chlamydia
  - Lacks muramic acid

## Cell Membrane

- Present in gram (+) and gram (-) bacteria
- Lipoprotein bilayer
- Electron transport and oxidative phosphorylation
- Enzymes and carrier molecules




Image courtesy Wikipedia/Public Domain

## Gram Positive Bacteria

- Unique feature: lipoteichoic acid (LTA)
- Major surface antigen for immune reaction
- In animal studies, LTA has induced:
  - Arthritis
  - Uveitis
  - Meningeal inflammation
  - Cascades resulting in septic shock and multi-organ failure
- Induces cytokine release
- Binds antibodies → activates complement cascade

## Gram Negative Bacteria

- Unique feature #1: Periplasm
  - Space between cell membrane and outer membrane
  - Contains many enzymes
  - B-lactamase → inactivates antibiotics




Image courtesy of Jeff Dahl/Wikipedia

## Gram Negative Bacteria

- Unique feature #2: Outer Membrane
  - Contains outer layer of lipopolysaccharide (LPS)
  - Major immune trigger for gram (-) bacteria





Image courtesy of Jeff Dahl/Wikipedia

## Lipopolysaccharide


- Components:
  - Polysaccharide
  - Lipid A
  - O antigen
- Lipid A
  - Highly toxic
  - Triggers cytokine release
- O antigen
  - Target for antibodies

## Key Point

- Different major surface antigens trigger the immune system in gram (+) and gram (-) bacteria
- Gram positive bacteria
  - Cell wall and membrane
  - Lipoteichoic acid (LTA)
- Gram negative bacteria
  - Outer membrane
  - Lipopolysaccharide (LPS)



## Capsules



## Capsules

- Sticky, gelatinous layer
- Secreted by bacteria
- Helps attach to host cells
- Protects against phagocytosis
- Mostly water with some polysaccharide
  - Special exception: Bacillus anthracis (anthrax)
  - Capsule is protein (d-glutamate)
  - Major virulence factor
  - Allows unimpeded growth

## Quellung Reaction


- Largely historical
- Used to detect strep pneumonia
- Rabbit antiserum added to bacterial slide
- Capsule swells when visualized under microscope
- Positive Quellung in encapsulated bugs
  - Strep pneumonia
  - H. influenza
  - N. meningitidis
  - E. Coli
  - Salmonella
  - Klebsiella
  - Group B strep (agalactiae)

## Capsules and Immunology

- B-cells secrete capsular antibodies (IgG)
  - Antibodies bind capsule
- Phagocytosis consume bacteria
  - Via Fc receptors
- Antibodies bind complement
  - Formation of MAC → cell death
  - Formation C3b → opsonin

## Capsules and Immunology

- Loss of antibodies/B-cells or complement
- Recurrent encapsulated bacterial infections



## Capsules and Immunology

- Asplenia
  - Risk of sepsis from encapsulated bacteria
  - Loss of splenic phagocytes

## Capsular Vaccines

- Capsular polysaccharides are basis for many vaccines
- Polysaccharides in capsule often weakly immunogenic
- “Conjugated” to an immune stimulator protein
  - diphtheria toxoid, tetanus toxoid, meningococcal outer membrane protein, mutant diphtheria protein
- Many conjugated vaccines for encapsulated bacteria
  - Neisseria meningitidis
  - Streptococcus pneumonia
  - Haemophilus influenzae type b

## Glycocalyx

- “Sugar coat” made of polysaccharides
- Similar to capsule
- Bacteria with distinct, firmly attached gelatinous layer have a capsule
- Bacteria with irregular, slimy fuzz layer have a glycocalyx
- Used to adhere to surfaces (i.e. catheters)
- S. epidermidis : biofilms




Image courtesy Wikipedia/Public Domain

## Pili and Fimbria

- Structurally similar to flagella
- Made of proteins
- Appendage or arm of bacteria
- Allows adherence to surfaces (ordinary pili)
- Attaches to another bacteria for conjugation (sex pili)
- Key bacteria:
  - E. Coli (UTIs/Pyelonephritis)
  - Neisseria Gonorrhoea (antigenic variation)

## Plasmids

- Small DNA molecule within a cell
- Physically separated from chromosomal DNA
- Can replicate independently
- Can contain genes for antibiotic resistance, toxins
- Can be transferred one bacteria to another




Image courtesy of Spaully/Wikipedia

## Flagellum

- Long, protein arms
- Used for motility

## Ribosomes

- Site of protein synthesis in bacteria
- Two subunits: 50S and 30S
  - S= sedimentation coefficient or Svedberg unit
- Different from ribosomes in eukaryotic cells
  - Allows selective toxicity of antibiotics
- Site of action of antibiotics
  - Tetracyclines: Bind to 30S subunit
  - Aminoglycosides: Interferes with 30S protein synthesis

## Spores

- Some bacteria can enter a dormant state called a spore
  - "Spore forming bacteria"
- Can survive long period of starvation
- Resistant to dehydration, heat, chemicals
- No metabolic activity

## Spores Components

- Coat:
  - Outermost layer
  - "Keratin-like" protein
  - Impermeable to many chemicals, antibacterial agents
- Cortex/Core Wall
  - Innermost layer
  - Peptidoglycans
- Dipicolinic acid
  - Large amounts inside spore
  - May help with heat resistance

## Spore Forming Bacteria

- *Bacillus anthracis*
- *Bacillus cereus*
- *Clostridium perfringens*
- *Clostridium tetani*
- *Clostridium Botulinum*

# Shapes and Stains


Jason Ryan, MD, MPH

## Identification of Bacteria


- Shape
- Color after staining
- Special tests

## Bacterial Shapes


- Coccus (sphere)
- Rod (bacilli)
- Coccobacillus
- Other



## Cocci



## Cocci



Streptococci  
"Chains"



Staphylococci  
"Bunches"  
"Clusters"

Image courtesy GrahamCoim/Wikipedia

Image courtesy Y Tambe/Wikipedia

## Cocci

- Most cocci are gram positive
  - Streptococcus
  - Staphylococcus
- Very few gram negative bugs are cocci
  - Neisseria (meningitidis/gonorrhoea)
  - Moraxella catarrhalis

## Rods

### Bacilli

- Most rods (and coccobacillus) are gram negative
- Few gram positive rods
  - Corynebacterium (diphtheria)
  - Clostridium
  - Listeria
  - Bacillus (anthrax, cereus)



Rod



Coccobacillus

## Other Shapes

- Branching/Filamentous
  - Resemble fungi
  - Actinomyces
  - Nocardia
- Spirochetes
  - Treponema (syphilis)
  - Borrelia (Lyme disease)
  - Leptospira (leptospirosis)
- Vibrio
  - Vibrio cholerae




Image courtesy Wikipedia/Public Domain

## Pleomorphic Bacteria

- Take on many shapes
- Rickettsia
- Chlamydia

## Common Bacterial Stains

- Gram Stain
- Giemsa
- Ziehl-Neelsen
- Silver
- India Ink – Cryptococcus (fungi)

## Simple Stains

- Methylene blue, safranin, and crystal violet
- Add to fixed preparation of bacteria
- Wash away
- Stain remains behind to show bacteria
- Used to see number bacteria, shapes

## Gram Stain





Image courtesy of Y tambe /Wikipedia



## Gram Stain

- Purple = Gram Positive
  - Retain crystal violet in cell walls
- Red = Gram Negative
  - Do not retain crystal violet in cell walls
  - Take up Safranin counter stain
- Thick cell wall of peptidoglycan in gram positive bacteria makes them purple

Image courtesy of Y tambe /Wikipedia

## Gram Stain Limitations

- Some bugs do not gram stain well
- Treponema (syphilis)
  - Too thin to see
- Mycobacteria (tuberculosis)
  - Mycolic acids in cell wall
- Mycoplasma
  - No cell wall
- Intracellular bacteria
  - Rickettsia (obligate intracellular)
  - Chlamydia (obligate intracellular; no muramic acid cell wall)
  - Legionella (mostly intracellular)

## Giemsa Stain

- Mixture of methylene blue, eosin, and Azure B
- Discoverer: Gustav Giemsa (1867-1948)
  - Looking for method to easily visualize plasmodium (malaria)
- Enters cells and stains nucleic acids
  - Used for blood smears, marrow

## Giemsa Stain

- Protozoa
  - Plasmodium
  - Trypanosomes
- Intracellular bugs
  - Chlamydia
  - Rickettsia
  - Borrelia (sometimes intracellular)

## Ziehl-Neelsen

- The "acid fast" stain
- Contains carbolfuchsin
- Used to detect mycobacterium (especially TB)
- Also used for Nocardia
- Acid fast bugs resists decolorization with acid solvents


Image courtesy Wikipedia/Public Domain

## Silver Stain

- Special stain for 3 organisms
- Pneumocystis pneumonia (HIV/AIDS)
  - Fungal infection
  - Diffuse interstitial pneumonia
- Legionella
  - Pneumonia
  - Contaminates water (outbreaks in nursing homes)
- H. Pylori
  - Gastric ulcers

## India Ink

- Negative stain
- Background stained, not bug
- Unstained organisms stand out in contrast
- Primarily used for *Cryptococcus neoformans*
  - Large polysaccharide capsule creates "halos"




India Ink stain shows yeast with "halos" Image courtesy of Crisco 1492

## Pigments


- Some bacteria produce special colors
- Staph Aureus
  - Golden, yellow color
- Pseudomonas aeruginosa
  - Blue-green pigment (pyocyanin)
- Serratia
  - Red pigment
- Actinomyces
  - Filamentous bacteria that "cement" together
  - Colonies have yellow-orange appearance
  - Known as "**sulfur granules**"

Pseudomonas (blue-green)



Wikipedia/Public Domain

Serratia (red)



Wikipedia/Public Domain

Staph Aureus (gold-yellow)



Matthias M./Wikipedia

# Bacterial Culture

Jason Ryan, MD, MPH

## Growth Plate

- Agar in Petri dish
  - Semi-solid substance from seaweed
  - Bacteria usually don't consume/decompose
- Nutrients added to support growth
  - Sugar
  - Water
  - Salts
  - Amino acids
- Many, many commercially available




Image courtesy Wikipedia/Public Domain

## Culture Media

- Non-selective
  - General purpose
  - Grows many bugs
  - Example: Nutrient agar
  - Also, blood agar: most commonly used non-selective media
- Selective
  - Contains toxic substances
  - Only certain bugs will grow
  - Thayer-Martin Media grows only Neisseria

## Culture Media

- Enriched
  - Special nutrients add so many bugs will grow
  - Blood agar
  - Chocolate agar
- Differential
  - Different bugs grow with different patterns
  - Blood agar: alpha, beta hemolysis

## Culture Media

- Blood agar
  - Enriched (blood)
  - Differential (hemolytic patterns)
- Eosin Methylene Blue
  - Selective (only gram negatives)
  - Differential (lactose fermenters)

## Fastidious Bacteria

- Fastidious = attentive to detail
- Fastidious organisms require special nutrients
- May not grow on standard media
- Some examples:
  - H. Influenza
  - Legionella

## Blood Agar


- Contain mammalian blood – usually 5% sheep blood
- Non-selective
- Enriched (blood)
- Differential by hemolysis pattern

**Hemolysis Patterns**  
 Beta = lysis  
 Alpha = partial  
 Gamma = no lysis



Image courtesy Y tambe/Wikipedia

## GRAM POSITIVE BACTERIA



## Blood Agar

- Commonly used to differentiate species of streptococcus
  - Alpha
  - Beta
  - Gamma
- Special feature of pseudomonas:
  - Beta-hemolytic
  - Greenish-metallic appearing colonies
  - Production of the pigments pyoverdinin and pyocyanin
- Staph Aureus
  - Beta hemolytic

## Chocolate Agar

- Variant of blood agar
- Contains red blood cells that are lysed (heating)
- Contains NAD (factor V) and hemin (factor X)
  - NAD from inside RBCs
  - Media heated such that they are not destroyed
- H. Influenzae will grow
- Classic scenario:
  - Bacteria won't grow on blood agar unless S. Aureus present


## Thayer-Martin Media

VPN/VCN

- Enriched, selective media for Neisseria
- Neisseria often from sites with lots of other flora
  - Throat, genitalia
  - Need very selective media
- Supplemented chocolate agar
- Vancomycin: Kills most Gram-positive organisms
- Colistin (polymyxin): Kills most Gram-negatives
  - Except Neisseria
- Nystatin: Kills most fungi

## Thayer-Martin Media

VPN/VCN



Wikipedia/Public Domain

## Bordet-Gengou Agar

Potato Agar

- Classic special media for *Bordetella pertussis*
  - Whooping cough
  - Extremely rare due to vaccination
- Prepared from potatoes → high instarch
  - Favorable to pertussis bacteria

## Loeffler's Media/Tellurite Plate

- Loeffler's Media
  - Selective media for *Corynebacterium diphtheriae*
- Tellurite Media (Cysteine-Tellurite Agar)
  - Differential media for *C. diphtheriae*
  - *C. diphtheriae* reduces potassium tellurite to tellurium
  - Produces gray-black colored colonies

## Lowenstein-Jensen Agar

- Special media for *Mycobacterium tuberculosis*
- Eggs, flour, glycerol, salt
- *M. tuberculosis* is SLOW growing
- Several weeks for visible colonies to appear
- *M. tuberculosis*: Ziehl-Neelsen stain

## Eaton's Agar

- Culture of *Mycoplasma pneumoniae*
- Bacteria has no cell wall
- Poorly visualized with gram stain
- Eaton medium specialized for *M. pneumoniae* growth
  - Require cholesterol to grow
- Takes days to weeks to grow
- Culture rarely used in modern era
- Diagnosis via:
  - Serology (antibody testing)
  - PCR (bacterial DNA)
  - Cold agglutinins (IM antibodies)
- Usually treated empirically


## MacConkey's Agar

- Selective media for gram (-) bacteria
- Contains bile salts as inhibitors of growth
- Inhibit Gram (+) bacteria
- Also differential for lactose fermenters
  - Lactose fermentation produces acid → turns agar pink
  - Non-lactose fermenters are colorless

## MacConkey's Agar



Image courtesy Wikipedia/Public Domain



### Eosin-Methylene Blue Agar

EMB

- Similar in function to MacConkey's Agar
- Eosin Y and methylene blue as inhibitors
- Inhibit Gram (+) bacteria
- Also differential for lactose fermenters
  - Lactose fermenters (Escherichia coli) appear as colonies with green metallic sheen or blue-black to brown color
  - Bacteria that do not ferment lactose appear as colorless or transparent colonies



### Sorbitol MacConkey Agar

- Detection of E. Coli O157:H7 strains (Shiga-like toxin)
- O157:H7 cannot ferment sorbitol (other E. Coli can)
- O157:H7 grows as colorless colonies on this medium
- Other E. Coli produce pink colonies

### Buffered Charcoal Yeast Extract

BCYE

- Contains dyes that give Legionella distinct color
- Antibiotics added: inhibits growth of competing bugs
- Very important to culture this bacteria
  - Can contaminate water supplies
  - Cause outbreaks
- Urinary antigen test also available
  - Only useful "type 1 infection"
  - Negative test is not 100%

### Sabouraud's Agar

- Selective media for fungi
- Developed by a French dermatologist
  - Growth of fungi in skin, hair, or nails (dermatophyte)
- Acid or antibiotics inhibit bacterial growth

# Special Growth Requirements

Jason Ryan, MD, MPH

## Bacterial Growth Environments

- Obligate anaerobes
- Obligate aerobes
- Facultative anaerobes
- Intracellular bacteria

## Energy Production

- Respiration
  - Electron transport chain
  - Makes ATP
  - Requires oxygen
- Fermentation
  - Sugars → acids
  - Makes ATP (less)
  - Does not use oxygen

## Superoxide Dismutase & Catalase

- Enzymes of aerobic organisms
- Superoxide radical ( $O_2^-$ ) produced by bacterial metabolism
- Superoxide dismutase
  - Catalyzes superoxide ( $O_2^-$ ) radical to  $O_2$  or hydrogen peroxide
- Catalase
  - Converts hydrogen peroxide ( $H_2O_2$ ) to oxygen and water
- Need these enzymes to survive in oxygen environments

## Obligate Aerobes

- Use  $O_2$  system to generate ATP
- Oxygen is final electron acceptor during respiration
- Can generate lots of energy (more than anaerobes)
- Contain superoxide dismutase
- Key bacteria:
  - *Pseudomonas aeruginosa*
  - *Mycobacterium tuberculosis*
  - *Nocardia* (opportunistic infections)

## Obligate Anaerobes

- Lack catalase or superoxide dismutase
- Common among normal flora of gut and mouth
  - 99% fecal flora
  - 100x more anaerobes than aerobes in mouth
- Don't cause communicable diseases
- Often live near mucosal surfaces
- Disease when surfaces breakdown
- Often present in abscesses
- Aminoglycosides ineffective (require  $O_2$ )

## Obligate Anaerobes

- Uses fermentation (no O<sub>2</sub>)
- Byproducts are often gases like CO<sub>2</sub> and H<sub>2</sub>
- Also produce short chain fatty acids
  - Acetic acid, isobutyric acid, many others
- Results in "foul smell"

## Obligate Anaerobes

- Actinomyces (gums; dental abscesses)
- Bacteroides (abdominal abscesses)
- Clostridium (botulinum; perfringens; tetani)

## Key Anaerobic Infections

- Abdominal abscesses/perforations
  - Contain many gram (-) flora of GI tract
  - Also contain Bacteroides fragilis (anaerobe)
  - B. fragilis resistant to many antibiotics
  - Treatment: Metronidazole + gram (-) agent
- Aspiration pneumonia
  - Mouth anaerobes enter lungs
  - Peptostreptococcus, Fusobacterium, Prevotella
  - Treatment: Clindamycin

## Facultative Anaerobes

- Can live without oxygen but use it if available
- Perform respiration and fermentation
- Pasteur effect: Oxygen inhibits fermentation
- Many common bacteria fall in this category
  - Staph
  - Strep
  - E. Coli

## Aerotolerant Anaerobes

- Similar to facultative anaerobes
- Always use fermentation even in presence of oxygen
- Rare
- Few examples relevant to clinical disease

## Obligate Intracellular Bacteria

- Cannot synthesize their own ATP
  - Depend on host for ATP
- Will not gram stain well (inside other cells)
- Difficult to grow (need cell culture)
- Rickettsia
  - Rocky Mountain spotted fever
  - Diagnosed clinically or with serology (antibody tests)
- Chlamydia
  - Diagnosis: Nucleic Acid Amplification Testing (DNA testing)

## Facultative Intracellular Bacteria

- Mycobacterium (macrophages)
- Legionella (macrophages)
- Salmonella (intestinal cells)
- Neisseria (urethral epithelial cells)
- Listeria (monocytes, macrophages)
- Brucella (macrophages and neutrophils)
- Francisella (macrophages)
- Yersinia pestis (macrophages)

# Virulence

Jason Ryan, MD, MPH

## Virulence Factors

- Bacterial features that allow evasion of host defenses
- Key examples to know:
  - Protein A
  - IgA protease
  - M protein

## Protein A

- Key virulence factor of Staph Aureus
- Part of peptidoglycan cell wall
- Inhibits phagocytosis
- Binds Fc portion of IgG antibodies
- Prevents opsonization and phagocytosis by macrophages
- Prevents complement activation

## IgA Protease

- Enzymes that cleave IgA
- IgA key for mucosal immunity
- Protease allows colonization of mucosal surfaces
- S. pneumonia
- H. influenza
- Neisseria (gonorrhoeae and meningitidis)

## M Protein

- Surface molecule of group A strep (pyogenes)
  - Strep throat, rheumatic fever
- M protein prevents phagocytosis
  - Binds factor H
  - Breaks down C3-convertase, prevent opsonization by C3b

## M Protein

- Shares properties with myosin
  - May be the basis of rheumatic heart disease
- Post-strep complications
  - Rheumatic heart disease
  - Glomerulonephritis
  - Different M protein subtypes associated each complication

## Bacterial Toxins

- Endotoxin
  - Only in gram (-) bacteria
  - Component of outer cell membrane
  - Lipopolysaccharide (LPS)
- Exotoxin
  - Proteins synthesized by some bacteria


## Endotoxin

- Component of gram (-) bacterial cell wall
- Released when bacteria die (not secreted)
- Lipopolysaccharide complex (LPS)
- Many different variants
- All have lipid A core
  - Responsible for most of the toxicity
- O antigen
  - Terminal end sugars that vary among bacterial strains
  - Do not cause disease by themselves

## Endotoxin

- Can cause fever, shock
- Triggers TNF and IL-1 release
  - Key immune components of sepsis and septic shock
- Generates weak antibody response
  - Can't vaccinate against endotoxin
- Heat stable (tolerates high temps)

## Endotoxin



## Lipooligosaccharide

LOS


- Endotoxin
- Similar to LPS with some structural differences
- Lacks O-antigen
- Found on non-enteric gram negatives
- Neisseria meningitidis is most important example

## Endotoxin

- Classic examples of endotoxin reactions
  - Meningococemia
  - Gram (-) sepsis

## Exotoxins

- Proteins secreted by bacteria → disease symptoms
- Classic structure: two component “A-B” polypeptide
- A component is toxic (A for active)
- B component binds to cell surfaces (B for binding)
- Various mechanisms of entry after B binding



## Exotoxins

- Many known exotoxins with various toxic effects
- General categories:
  - Inhibit protein synthesis
  - Increase fluid secretion
  - Inhibit phagocytosis
  - Inhibit neurotransmitter release
  - Lyse cell membranes
  - Superantigens

## Toxin Mechanisms

### Protein Synthesis Inhibitors


- *Corynebacterium diphtheria*
- *Pseudomonas aeruginosa*
- *Shigella*
- Enterohemorrhagic *E. Coli* (EHEC)

## ADP Ribosylation

- Two toxins work by adding ADP-Ribose to proteins
  - Diphtheria toxin
  - Endotoxin A (*pseudomonas aeruginosa*)
- Addition of ADP-Ribose makes protein dysfunctional


## *Corynebacterium Diphtheria*

- Sore throat with membrane, swollen nodes
- Largely eradicated by vaccination
  - Diphtheria-pertussis-tetanus (DPT) vaccine
- Diphtheria toxin: Inactivates elongation factor (EF-2)
- EF-2 necessary for protein synthesis
- Lethal toxin



## *Pseudomonas aeruginosa*

- Causes many types of infection
  - Skin, sepsis, pneumonia
- Secretes exotoxin A
- Same mechanism as diphtheria toxin



## Shigella

- Causes infectious diarrhea
- Secretes shiga toxin
- Binds to 60S ribosome in cells
- Removes a specific adenine residue from rRNA in the 60S ribosomal subunit
- Halts protein synthesis
- Special note:
  - Invasion of GI mucosal cells is main cause of disease
  - Non-toxicogenic strains cause significant disease

## Enterohemorrhagic E. Coli (EHEC)

- Some E. Coli strains produce “shiga-like” toxin
- Same mechanism as shiga toxin
- Typically causes bloody diarrhea
- Classic serotype is E. coli O157:H7
- Do not invade host cells
  - Attach to intestinal epithelial cells
  - Disease from secretion of proteins into host cells
  - Toxin

## Shiga Toxin

- Also stimulates cytokine release
- When reaches systemic circulation, can lead to hemolytic uremic syndrome

## Key Points


- Shigella and EHEC produce shiga toxins
- Both cause bloody diarrhea
- Shigella
  - Disease from bacterial invasion of mucosa
  - Toxin less important in disease than invasion
- EHEC
  - Do not invade cells
  - Disease from toxin (inflammation)
  - Hemolytic uremic syndrome

## Toxin Mechanisms

Increase Fluid Secretion

- Enterotoxigenic E. Coli (ETEC)
- Bacillus anthracis
- Vibrio cholera

## Fluid Secretion in GI Tract



Key Points:

- #1: Amount of Cl<sup>-</sup> secreted ≈ amount of water in GI tract
- #2: To increase Cl<sup>-</sup> secretion, activate Gs or AC

## Enterotoxigenic E. Coli

(ETEC)

- Two toxins differentiated by heat stability
  - Heat labile toxin (LT)
  - Heat stable toxin (ST)

## E. Coli Heat Labile Toxin

- Activates adenylate cyclase ( $\uparrow$ cAMP)
- Increases water in gut  $\rightarrow$  diarrhea

## E. Coli Heat Stable Toxin

- Activates guanylate cyclase
- Increases cGMP
- Stimulation of chloride secretion
- Inhibition of sodium chloride absorption
- More water in gut  $\rightarrow$  diarrhea

## Bacillus Anthracis

- Anthrax makes three proteins: protective antigen (PA), lethal factor (LF), and edema factor (EF)
- Alone they cause no known physiological effects
- In pairs they produce toxicity
- Edema toxin = PA + EF
- Mimics adenylate cyclase
- Multiple sites of disease
  - Skin (most common)
  - Lungs (inhalation  $\rightarrow$  necrotizing pneumonia)
  - GI (ulcers)
- Skin and GI lesions often have edematous borders
  - May be caused by edema factor

## Vibrio Cholera

- Cholera toxin
- Permanently activates  $G_s \rightarrow \uparrow$ cAMP
- Voluminous "rice-water" diarrhea
- Common in areas with lack of clean water
- Death: profound dehydration, electrolyte losses, shock
- Aggressive volume repletion is mainstay of treatment

## Toxin Mechanisms

Inhibitors of Phagocytosis

- *Bordetella pertussis* (whooping cough)
- Pertussis toxin
- Shown to inhibit  $G_i$  proteins
- Allows over-activation of adenylate cyclase
- $\uparrow$ cAMP levels in cells in neutrophils
- Result: impaired recruitment of neutrophils


## Toxin Mechanisms

### Neurotoxins

- Clostridium tetani
- Clostridium botulinum
- Both work by disruption of SNARE proteins

## SNARE Proteins

SNARE proteins present in vesicles  
Allow vesicles to "dock" and unload NT



## Neurotoxins

- Clostridium tetani
  - Tetanospasmin
  - Works in spinal cord (Renshaw cells)
  - Inhibits inhibitory neurons (GABA and glycine)
  - Result: Muscles always on (rigid)
- Clostridium botulinum
  - Botulinum toxin
  - Works at neuromuscular junctions
  - Prevents Ach release (no muscle contraction)
  - Result: Muscles floppy (flaccid paralysis)

## Toxin Mechanisms

### Lysis of Cell Membranes

- Clostridium perfringens
- Streptococcus pyogenes

## Clostridium perfringens

### Gas gangrene

- Alpha toxin
- Phospholipase C enzyme
- Degrades phosphatidylcholine and sphingomyelin
- Muscle breakdown (myonecrosis)
- Causes a decline in muscle blood flow
- Forms occlusive plugs: platelets, leukocytes, fibrin
- Result: Low O<sub>2</sub> environment favorable to bacteria

## Strep Pyogenes

### Strep Throat/Rheumatic Fever /Glomerulonephritis

- Streptolysin O
- "Cytolysin" (lysis cells)
- Responsible for beta hemolysis (also streptolysin S)
- Anti-streptolysin O (ASO) antibodies
  - Elevated following strep infection
  - Can be useful in suspected rheumatic heart disease or post-strep glomerulonephritis

## Toxoid Vaccines


- Toxoid = inactivated bacterial toxin
- Used for vaccination
- Used to prevent diphtheria and tetanus
- Part of DTaP combined immunization
  - Diphtheria
  - Tetanus
  - "Acellular" Pertussis (inactive toxin plus bacterial elements)

## Exotoxin Genetics

- Many exotoxin genes not part of chromosome
- Plasmid-encoded
  - E. coli heat-labile toxin
- Bacteriophage-encoded
  - Corynebacterium diphtheriae
  - Strep pyogenes erythrogenic
  - E. Coli shiga-like toxin
  - Botulinum toxin
  - Cholera toxin

## Superantigens

- Activate a MASSIVE number of T-cells



## Superantigens

- Typical antigen response: <1% T-cells
- Superantigen: 2-20% T-cells
- HUGE release of cytokines (IFN- $\gamma$  and IL-2)
- Massive vasodilation and shock

## Superantigens

- Staph aureus
  - Toxic shock syndrome toxin (TSST-1)
- Strep pyogenes (group A strep)
  - Pyrogenic exotoxin A or C
- Both can cause toxic shock syndrome
  - More common with staph
  - Strep TSS often associated with necrotizing fasciitis


## Toxic Shock Syndrome

- Fever, shock (hypotension), red rash
- Diffuse, red erythroderma (resembles sunburn)
- After weeks: desquamation of palms/soles
- Diarrhea is common
- Multi-organ system failure often results
- Classic scenarios (staph + packing):
  - Women using tampons
  - Surgical wound with packing


# Growth and Genetics

Jason Ryan, MD, MPH


## Bacterial Growth Curve




## Bacterial Growth Curve




## Bacterial Growth Curve




## Bacterial Growth Curve



## Bacterial Growth Curve



## Bacterial Growth Curve



## Gene Transfer

- Bacteria often transfer genetic material
- Key for evolution of antibiotic resistance
- Three key mechanisms:
  - Transformation
  - Conjugation
  - Transduction

## Bacterial Transformation

- Direct uptake DNA from **surrounding environment**
- Allows for evolution of DNA over time
- Very useful technique in micro labs
- Introduce genes to bacteria for replication

## Bacterial Conjugation

- Transfer from one cell to another **via pilus**
  - Physical contact of two organisms
- DNA transferred **via plasmids**

## Plasmids

- Small DNA molecule within a cell
- Physically separated from chromosomal DNA
- Can replicate independently
- Can contain genes for antibiotic resistance, toxins
- Can be transferred one bacteria to another




Image courtesy of Spaully/Wikipedia

## Bacterial Conjugation





Image courtesy of Adenosine/Wikipedia

## Transformation vs. Conjugation

- Classic scenario:
  - Strain A requires amino acid X for growth (does not need Y)
  - Strain B requires amino acid Y for growth (does not need X)
  - Strain A and Strain B grown together without X or Y
  - DNAase added to medium to degrade DNA
  - Bacteria grow!
  - Ability to grow without X/Y xferred between bacteria
  - Cannot be transformation
    - DNAase destroyed any leaked DNA
  - Must be conjugation

## High Frequency Strains



## High Frequency Strains

- Used to map genes
- Process takes time
- Can interrupt at various time intervals
- See which genetic material has transferred
  - Plasmid site is origin of genetic transfer
  - Initial material transferred is that closest to plasmid
- With multiple experiments can make a map

## Transduction

- Transfer of DNA via a **bacteriophage**
  - Virus that infects bacteria
- Virus picks up DNA, transfers to another bacteria

## Transduction

- Generalized
  - Virus infects bacteria
  - Multiplies, randomly picks up host DNA
  - Host DNA transferred to other bacteria




Image courtesy of Wikipedia/Public Domain

## Transduction

- Specialized
  - Transfer of specific genes
  - Virus DNA inserts into host DNA (lysogeny)
  - When bacteriophage DNA excised, packaged into virus with specific host DNA

## Lytic vs. Lysogenic Phages

- Transduction happens in two ways
- Lytic cycle
  - Nuclear material enters bacteria
  - Multiplies, lyses cell
  - Releases progeny viruses
- Lysogenic cycle
  - Nuclear material enters cell
  - Incorporates in host DNA
  - May later become excised (enter lytic phase)
- Phages that replicate only via the lytic cycle: virulent
- Phages that replicate using both: temperate

## Lytic Phages

- Virus infects bacteria
- Uses cellular machinery to reproduce
- Lyses cell
- Usually generalized gene transfer





Image courtesy of Wikipedia/Public Domain

## Lysogenic Phages

- Virus infects bacteria
- Incorporates phage DNA into bacterial DNA
- Can remain dormant for long periods of time
- Certain triggers (i.e. UV light) induce genome excision
- Results in a lytic cycle and release of phage particles



Asiela/Wikipedia

## Why Lysogeny Matters

- Genes for some **bacterial toxins** are transferred to non-toxic strains via lysogeny
- Example:
  - Not all strains *C. diphtheria* are toxic
  - Gene for toxin is not part of bacteria's genome
  - Gene carried by a phage (corynephage)
  - Toxicity depends on infection with phage

## Lysogenic Toxins

- Diphtheria toxin
- Erythrogenic toxin (*S. pyogenes*; Scarlet fever)
- Shiga-like toxin (*E. Coli*; EHEC)
- Cholera toxin
- Botulinum toxin

Source: Brussow, et al. Microbiol Mol Biol Rev. Sep 2004; 68(3): 560-602


## Transposition

- Transposons are DNA segments within bacterial DNA
- Can be excised and reintegrated in new locations in DNA
- Once excised, can also be moved to plasmid
- Mechanism of transfer of **resistance to antibiotics**
  - Bacteria #1 is resistant
  - Transposon segment carries resistance gene
  - Transposon moved to plasmid
  - Plasmid transferred to other bacteria

# Bacterial Identification

Jason Ryan, MD, MPH


## Gram Positive Bacteria




## Lancefield Grouping

- System for classifying streptococci
- Based on "C carbohydrates" in the cell wall that allow agglutination with particular antisera
- Commercially available tests for different antigens
- Clinically relevant groups:
  - Group A: Strep Pyogenes
  - Group B: Strep Agalactiae
  - Group D: Enterococcus

## Cocci



## Cocci



Streptococci  
"Chains"



Staphylococci  
"Bunches"  
"Clusters"

Image courtesy GrahamCoim/Wikipedia

Image courtesy Y Tambe/Wikipedia

## Blood Agar

- Contain mammalian blood - usually 5% sheep blood
- Non-selective
- Enriched (blood)
- Differential by hemolysis pattern


**Hemolysis Patterns**  
Beta = lysis  
Alpha = partial  
Gamma = no lysis



Image courtesy Y tambe/Wikipedia

## Catalase Test

- Differentiates Staph from Strep
- Catalase enzyme produced by bacteria that use oxygen
- Breaks down  $H_2O_2$  into  $H_2O$  and  $O_2$  (makes bubbles)
- Hydrogen peroxide on slide
- Add bacteria
- Look for bubbles
- Catalase positive = bubbling
- Catalase negative = no bubbling



CDC/Public Domain

## CGD

Chronic Granulomatous Disease

- Phagocytes use NADPH oxidase
- Generate  $H_2O_2$  from oxygen (respiratory burst)
- CGD = Loss of function of NADPH oxidase
  - Phagocytes cannot generate  $H_2O_2$
- Catalase (-) bacteria generate their own  $H_2O_2$  which phagocytes use despite enzyme deficiency
- Catalase (+) bacteria breakdown  $H_2O_2$ 
  - Host cells have no  $H_2O_2$  to use → recurrent infections
- Five organisms cause almost all CGD infections:
  - Staph aureus, Pseudomonas, Serratia, Nocardia, Aspergillus

Source: UpToDate

## Coagulase Test

- Differentiates Staph Aureus from other Staph
- Rabbit plasma in tube, add bacteria
- Coagulase (cell surface) causes fibrin clot to form
- Coagulase positive = clumping
- Coagulase negative = no clumping

## Novobiocin

- Differentiates *S. saprophyticus* from *S. epidermidis*
- Technique:
  - Plate bacteria on agar with Novobiocin "disk"
  - Measure clearance zone around disk
- Resistant = Growth near edge of disk
- Sensitive = Large zone of clearance around disk



CDC/Public Domain

## Bacitracin

- Differentiates Group A strep from Group B strep
- Bacitracin: antibiotic that interferes with peptidoglycan synthesis
- Bacteria vary in their susceptibility
- Technique:
  - Plate bacteria on agar with bacitracin disk
  - Measure clearance zone around disk
- Resistant = Growth near edge of disk
- Sensitive = Large zone of clearance around disk

## Optochin

ethylhydrocupreine

- Differentiates *S. pneumoniae* from *S. viridans* strep
- *S. pneumoniae* highly sensitive to Optochin
- Technique:
  - Plate bacteria on agar with optochin disk
  - Measure clearance zone around disk
- Resistant = Growth near edge of disk
- Sensitive = Large zone of clearance around disk


## Bile Esculin Agar

- Selective for:
  - Group D strep (Enterococci and *S. bovis*)
- Bile salts inhibit most Gram-positive bacteria
- Esculin:
  - Hydrolyzed by Group D strep
  - Media turns dark brown/black

## NaCl Media

- Differentiates Enterococcus from non-enterococcus Group D bacteria (*S. bovis*)
- Enterococcus is "salt tolerant"
- Can grow in high salt concentrations
- Inoculate bacteria on high NaCl media
- Watch for growth

## Lancefield Group D




## Mannitol Salt Agar



- Used to differentiate staph species
- Contains high concentration of salt (7.5%)
- Staph can tolerate high saline levels
- Contains mannitol and pH indicator (phenol red)
- Mannitol fermenters make acid
- Phenol red turns yellow
- *Staph aureus* ferments mannitol
- Most other staph do not
- **Growth on MSA with yellow color is *Staph Aureus***

## Gram Negative Bacteria



## Maltose

- *Neisseria meningitidis* can metabolize maltose
- *Neisseria gonorrhoeae* cannot
- Growth media with maltose used to differentiate

## Lactose Fermentation

- MacConkey's Agar
- Selective media for gram (-) bacteria
- Contains bile salts as inhibitors of growth
- Inhibit Gram (+) bacteria
- Inhibits fastidious gram (-): Neisseria, Pasteurella
- Differential for lactose fermenters
  - Lactose fermentation produces acid → turns agar pink
  - Non-lactose fermenters are colorless

## MacConkey's Agar



Image courtesy Wikipedia/Public Domain

## Fast and Slow Fermenters

- Citrobacter and Serratia can initially appear as non-lactose fermenting due to slow growth
- Longer incubation will show growth

## Oxidase

- Test for presence of cytochrome oxidase
- Bacterial colonies placed on paper discs with indicator present
- If oxidase is present, color change occurs

## Campylobacter & Vibrio


- Oxidase (+) organisms that are "comma shaped" may be Campylobacter or Vibrio
- Campylobacter grows at 42°C (Vibrio does not)
- Vibrio grows on alkaline media (Campy does not)

## H<sub>2</sub>S Production

- Oxidase (-) organisms can be subdivided by H<sub>2</sub>S
- Salmonella and proteus produce H<sub>2</sub>S
- Shigella does NOT produce H<sub>2</sub>S
- Triple Sugar Iron (TSI) test
  - Organisms that produce H<sub>2</sub>S will turn TSI media black

# Staphylococci

Jason Ryan, MD, MPH



# Staphylococci

- Staph Aureus
- Staph Epidermidis
- Staph Saprophyticus
- All gram positive cocci
- All form clusters
- All catalase (+)

# Staph Aureus

## Special Features

1. Basic habitat is the nares (nose)
  - 30% of people carry the bacteria
2. Produces a yellow pigment (aureus = golden)
3. Beta-hemolytic
4. Coagulase (+)
  - Forms fibrin clot
5. Protein A virulence factor
  - Blocks Fe-IgG interaction
  - Prevents phagocytosis & complement activation
6. Produces several toxin-related diseases

# Staph Aureus

## Infections

- Toxin disease
  - Toxic Shock Syndrome (TSST-1)
  - Food poisoning (Staph aureus enterotoxin)
  - Scalded skin syndrome (Exfoliatin)
- Infectious diseases
  - Skin infections (Impetigo)
  - Pneumonia
  - Endocarditis
  - Osteomyelitis
  - Abscesses

# Toxic Shock Syndrome

- Toxic Shock Syndrome Toxin (TSST-1)
- Fever, shock (hypotension), red rash
- Diffuse, red erythroderma (resembles sunburn)
- After weeks: desquamation of palms/soles
- Diarrhea is common
- Multi-organ system failure often results
- Classic scenarios (staph + packing):
  - Women using tampons
  - Surgical wound with packing


## Staph Food Poisoning

- Food handler contaminates food
- Food left at room temperature several hours
  - Picnic is classic scenario
- Bacteria grow in food → produce enterotoxin
- Ingestion of preformed toxin causes disease
- GI illness develops ~3 to 6 hours later
  - Nausea, vomiting (diarrhea rare)
  - Abdominal cramps
- Look for multiple sick people after eating at a picnic
- Classic food is mayonnaise in potato or egg salad

## Scalded Skin Syndrome

- Newborn disease
- Caused by *S. Aureus* exfoliative toxin (Exfoliatin)
- Classically occurs 3 to 7 days of age
- Fever, diffuse erythema starting at the mouth
- Sloughing of skin
- Toxin destroys keratinocyte attachments in stratum granulosum only
- Damage intraepidermal: Heals completely (no scar)
- Nikolsky's sign: skin slips off with gentle tug

## Skin Layers



Mikael Haggström, /Wikipedia

## Bullous Impetigo

- Impetigo = skin infection
  - Caused by Group A strep or Staph Aureus
  - Honey colored, crusted lesions
- Bullous impetigo = variant of impetigo with bullae
- Bullae = fluid-filled sacs similar to blisters
- Bullous impetigo caused by *S. Aureus*
- Classically occurs in children
- Easily spread one child to another
- *S. Aureus* exfoliative toxin strains



Image courtesy of Åsa Thörn/Wikipedia

## Pneumonia

- Staph is rare cause of lobar pneumonia
- Classically occurs as "post-infectious"
  - Bacterial pneumonia following influenza

## Endocarditis

- Classic cause of ACUTE endocarditis
  - Rapid onset of symptoms
  - Very ill patient
  - Often no pre-existing valve disease (i.e. mitral valve prolapse)
- Contrast with subacute → Strep Viridans
  - Slower onset of symptoms
  - Less sick patient
  - Prior valve abnormality

## Intravascular devices

- “Central lines”
- Common cause of staph bacteremia
- Most important preventative measure is sterile technique:
  - Wash hands
  - Gloves
  - Sterile insertion practices

## Osteomyelitis

- S. Aureus is common cause of osteomyelitis
- Children: Usually long bones (femur, tibia, fibula)
- Adults: Usually spine
- Mechanisms:
  - Hematogenous spread
  - Spread from skin/soft tissues
  - Trauma (surgery)
- Symptoms usually localized pain +/- fever
- Diagnosis made by imaging (CXR, CT scan, MRI)

## Osteomyelitis

### Classic Causes

- Child
  - Staph aureus (hematogenous spread)
- Sickle Cell patient
  - Salmonella (hematogenous spread)
- TB patient
  - Pott's disease (vertebrae/spine)
- Diabetic
  - Polymicrobial from foot ulcer
- Bedbound patients
  - Polymicrobial from pressure sores

## Cellulitis

- Infection of deep dermis and subcutaneous fat
- Mostly caused by  $\beta$ -hemolytic streptococci
- S. Aureus can also cause
- Antibiotics must cover Staph



Image courtesy of Pshawnosh/Wikipedia

## Abscesses

- Bacteria and inflammatory cells (pus)
- Walled off in deep tissues
- Skin abscesses commonly caused by S. Aureus
  - Furuncle = boil; infection of hair follicle
  - Carbuncle = multiple boils clustered together
- Tonsillar abscesses
- Mainstay of treatment is incision and drainage

## Staph Aureus Antibiotics

- Most strains of Staph resistant to penicillin
- Produce beta-lactamases
- Antistaphylococcal penicillins
  - Dicloxacillin, Nafcillin, Oxacillin
- First generation cephalosporins
  - Cephalexin
- Beta lactam plus inhibitor
  - Amoxicillin/clavulanate

## MRSA

Methicillin-resistant *Staphylococcus aureus*

- Resistant to all beta lactams
- Altered penicillin binding proteins (PBPs)
- Important hospital-acquired bacteria
  - Sometimes community acquired
- Vancomycin or daptomycin: antibiotics of choice
- Can also use Linezolid

## Staph Epidermidis

- Normal skin flora; two clinical implications
- #1: Blood culture contaminant
  - Needle/IV contaminated by *S. epi*
- #2: Infects prosthetic materials in blood
  - Surface molecules aid in adherence
  - Bacteria produce biofilms
  - Catheter infections
  - Pacemaker infections
  - Prosthetic heart valves
  - Prosthetic joints
- Often methicillin resistant
- Treatment: Vancomycin

## Staph Epidermidis

- Often methicillin resistant
  - Resistance to methicillin >80 percent
- Treatment: Vancomycin


## Staph Saprophyticus

- Cause of UTIs especially in sexually active women
  - Most caused by *E. Coli* (~90%)
  - Other bugs: *Proteus*, *Klebsiella*, *S. Saprophyticus*
- Key features:
  - Sexual activity (honeymooner's cystitis)
  - Nitrite negative on urine dipstick
- Treated by UTI antibiotics
  - Fluoroquinolones
  - SMX-TMP
  - Nitrofurantoin

# Streptococci

Jason Ryan, MD, MPH

## Gram Positive Bacteria



## Streptococci

- Strep Pyogenes (Group A)
- Strep Agalactiae (Group B)
- Strep Pneumoniae
- Strep Viridans
- Enterococcus (Group D)
- Strep Bovis
- All gram positive cocci
- All form chains
- All catalase (-)

## Strep Pyogenes

Group A Strep

- Pyrrolidonyl arylamidase (PYR) positive
  - Substrate (L-naphthylamide-β-naphthylamide) hydrolyzed by bacterial enzyme to β-naphthylamide
  - Can be detected by color change with detection reagent
- M protein virulence factor
  - Inhibits phagocytosis

## Strep Pyogenes

Group A Strep

- Cause of many different illnesses
- Infections
  - Pharyngitis (Strep throat)
  - Skin: Cellulitis/Impetigo
- Toxin-mediated disease
  - Scarlet fever
  - Necrotizing fasciitis
  - Toxic Shock Syndrome
- Immune disease
  - Rheumatic fever
  - Post-strep glomerulonephritis

## Strep Pharyngitis

- 15-30% pharyngitis due to S. pyogenes
  - Many cases viral
- Important to identify and treat S. pyogenes
  - Prevent disease transmission
  - Limit symptoms, severity
  - Prevent rheumatic fever
- Diagnosis:
  - Throat culture
  - Rapid antigen test (useful if positive)
- Treatment: Penicillin, amoxicillin, cephalosporins

## S. Pyogenes Skin Infections

- Cellulitis and Impetigo
- Both commonly caused by Strep but also S. Aureus
- Antibiotics need to cover Strep and Staph



Image courtesy of Pshawnoah/Wikipedia



Image courtesy of Åsa Thörn/Wikipedia

## Scarlet Fever

- Rash following pharyngitis
- Skin reaction to erythrogenic toxin
- Gene for toxin transferred by lysogenic phage

## Scarlet Fever

- Fever, sore throat, diffuse red rash
- Also, many small papules ("sandpaper" skin)
- Starts head/neck → expands to cover trunk
- Classic finding: Strawberry tongue
- Eventually skin desquamates
- Palms and soles are usually spared



Image courtesy of Wikipedia/Public Domain



Image courtesy of Afag Azizova/Wikipedia

## Necrotizing Fasciitis

- Infection of deep tissues
  - Muscle fascia/subcutaneous fat
  - Streptococcal pyrogenic exotoxin released
- Often fulminant and deadly
  - Infection spreads along muscle fascia
- Requires urgent surgical debridement

## Necrotizing Fasciitis


- Type 1:
  - Polymicrobial
  - Often anaerobes (Bacteroides, Clostridium, etc.)
  - Occurs in diabetics, immunocompromised, post surgery
- Type 2:
  - Group A strep (sometimes Staph)
  - Occurs in otherwise healthy people after skin injury
- Classic case:
  - Minor skin trauma
  - Redness/warmth (can be confused with cellulitis)
  - Pain out of proportion to exam
  - Fever, hypotension

## Rheumatic Fever


- Follows group A strep pharyngitis infection
- Streptococcus → anti-strep antibodies
- Cross react with tissue antigens
- Type II autoimmune reaction

## Rheumatic Fever

- Aschoff bodies
  - Cardiac nodules with inflammatory cells (granulomas)
  - Pathognomonic for rheumatic carditis
- Anitschkow's cells
  - Macrophages with owl eye appearance
- Elevated ASO titers



Nephron/Wikipedia



Ed Uthman, MD/Wikipedia

## Jones Criteria

Major	Minor
Carditis	Fever
Polyarthritits	Arthralgia
Chorea	Prior RF
Erythema marginatum	↑WBC, ESR, CRP
Subcutaneous nodules	Prolong PR interval

2 Major or 1 Major & 2 Minor  
(Must have evidence of strep infection )

## Rheumatic Fever

- Clues:
  - Sore throat or URI followed by joint pain, new murmur
- Treatment: Penicillin
- Cardiac involvement
  - Acute RF → severe valve disease, heart failure
  - Later → mitral stenosis (rarely aortic or tricuspid valves)

## Post-streptococcal GN

- Nephritic syndrome 2-3 weeks after GAS infection
- Nephritogenic strains
  - Bacteria with certain M protein subtypes cause nephritis

## Strep Agalactiae

Group B Strep

- Beta hemolytic (like GAS)
- Makes CAMP factor
  - Staph Aureus makes  $\beta$ -hemolysin
  - CAMP factor enhances lysis by  $\beta$ -hemolysin
- Hydrolyzes the compound hippurate
  - Will alter color of hippurate test




Image courtesy of Bluetritidum/Wikipedia

## Strep Agalactiae

Group B Strep

- Colonizes vagina
- Causes infections in newborns
  - Babies infected in utero or during birth
  - Pneumonia, meningitis, sepsis
- Pregnant women screened 35-37 weeks
  - Vaginal culture
- Women GBS (+) receive prophylaxis
  - Four hours prior to delivery
  - Penicillin, ampicillin, or cefazolin

## Strep Pneumonia

- “Lancet shaped” gram (+) cocci
- IgA protease
- Key virulence factor: Polysaccharide capsule
  - Prevents phagocytosis
  - Basis of vaccine from capsular material
  - Asplenic patients at risk for sepsis
    - Splenectomy
    - Sickle cell anemia

## Strep Pneumonia

- Lobar Pneumonia
- Meningitis
- Otitis Media
- Sinusitis

## Viridans Group Strep

- Group of similar bacteria
- No Lancefield group (A, B, D)
- Normal mouth flora
- Cause dental carries (Strep Mutans)
- SUBACUTE endocarditis (Strep Sanguis)
  - Slow onset symptoms; less sick patient
  - Often affects ABNORMAL valves
  - Dextran → fibrin
  - Requires endothelial damage
  - Mitral valve prolapse
  - Pearl: Recent dental procedure

## Enterococcus

- E. faecalis, E. faecium
- Normal colonic bacteria
- Lancefield group D
- Infections:
  - UTIs
  - Endocarditis (rare)
- Resistant to penicillin
- Vancomycin resistant enterococcus (VRE)
  - Dangerous hospital acquired infection

## Enterococcus

- Relatively resistant to cell wall agents
  - Penicillin, ampicillin, and vancomycin
- Impermeable to aminoglycosides
- Bacteremia: Often use synergistic therapy
  - Vancomycin/gentamycin
  - Ampicillin/gentamycin
- Vancomycin resistant enterococcus (VRE)
  - Dangerous hospital acquired infection
  - Linezolid, Daptomycin
- Micro lab sensitivities very important for therapy

## Enterococcus UTIs

- Often hospital acquired, associated with catheters
- Removal of catheter alone may cure infection
- Urinalysis: NEGATIVE for nitrites

## Enterococcus Endocarditis

- Rare cause of endocarditis
- Usually after manipulation GU tract
  - TURP procedure, cystoscopy
- Vancomycin/gentamycin often used for empiric therapy before culture data available
- Synergistic effect of dual antibiotics


## Strep Bovis

- Normal colonic bacteria
- Rare cause bacteremia/endocarditis
- Strongly associated with colon cancer

## Other Gram Positives

Jason Ryan, MD, MPH

## Gram Positive Bacteria



## Clostridium

- Family of bacteria
  - All form spores
  - All obligate anaerobes
  - All form toxins
- *C. tetani* (tetanus)
- *C. botulinum* (botulism)
- *C. perfringens* (wound infections, food poisoning)
- *C. difficile* (diarrhea after antibiotic use)

## Clostridium tetani

- Spores found in soil
- Enter body via penetrating injury
  - Classic scenario: barefoot on rusty/dirty nail or splinter
- Spores germinate into bacteria
- Tetanus toxin produced (tetanospasmin)
- Travels to spinal cord
- Blocks glycine and GABA release by inhibitory neurons
  - "Renshaw cells:" inhibitory spinal cord interneurons

## Clostridium tetani

- Spasms, muscle contractions, rigidity
- Classic symptoms
  - Lockjaw (trismus)
  - Risus sardonicus (forced grin due to spastic facial muscles)

## Clostridium tetani

- Treatment for tetanus
  - Wound debridement
  - Metronidazole
  - Tetanus immune globulin (binds circulating toxin)
  - Benzos or neuromuscular blockers until toxin wears off
- Tetanus toxoid used for vaccination

## Clostridium botulinum

- Ubiquitous organisms
- Vegetables, fruits, seafood, soil
- Heat-resistant spores
  - Survive up to 100°C
- Botulinum toxin
  - Works at neuromuscular junctions
  - Prevents Ach release (no muscle contraction)
  - Result: Floppy muscles (flaccid paralysis)
- Many different variants of toxin
- Some carried by bacteriophages

## Botulism

- Three types: food, infant, wound
- Food (toxin ingestion; usually adults)
  - Undercooked food
  - Canned food: anaerobic environment promotes growth
  - Watch for multiple sick adults after a meal
- Infant (spores)
  - Ingestion of spores → growth in infant intestine
  - Watch for contaminated honey!
- Wound (bacterial growth)
  - Infection with *C. botulinum*

## Botulism

- Symptoms: 12-48 hours after ingestion
- Symptoms: 3 D's
  - Diplopia, dysphagia, dysphonia
- Diagnosis:
  - Often clinical
  - Spores and toxins sometimes detected in stool
- Treatment:
  - Antitoxin blocks circulating toxin
  - Cannot block toxin already in nerves
  - Supportive care → toxin washout

## Clostridium perfringens

- Widespread in nature, especially soil
- Infects dirty wounds & causes food poisoning
- Causes gas gangrene (clostridial myonecrosis)
  - Traumatic wound with vascular compromise
  - Favorable environment for anaerobic growth
- Alpha toxin
  - Destroys muscle tissue and causes hemolysis
  - Phospholipase that acts on lecithin (lecithinase)
  - Degrades phospholipids in cell membranes

## Gas Gangrene

- Severe pain at injury site within 24 hours
- Skin tense and tender
- Systemic toxicity
  - Fever, Hypotension, Shock
- Diagnosis
  - Gas at injury site on imaging
  - Crepitus
- Treatment
  - Surgical debridement
  - Broad spectrum antibiotics



Image courtesy of Engelbert Schröpfer, Stephan Rauthe and Thomas Meyer/Wikipedia

## Clostridium perfringens

- Food poisoning (undercooked meats)
  - Spores ingested → produce toxin
  - Late onset (8-22hrs) watery diarrhea
  - Contrast with *S. aureus*/*B. cereus* (preformed toxin)

## Clostridium difficile

- Ubiquitous spores in nature including soil
- Ingestion not harmful with normal GI flora
  - Colonic flora prevent overgrowth of C. diff
- Causes antibiotic-associated colitis
  - Antibiotics alter normal gut flora
  - Favorable environment for C. diff growth

## Clostridium difficile

- Not invasive: disease via toxins
- Two toxins
  - Toxin A: Enterotoxin → watery diarrhea
  - Toxin B: Cytotoxin → Cell necrosis/fibrin deposition
  - Both bind to GI cells and are internalized
  - Destroy cytoskeleton of GI cells → pseudomembrane

## Clostridium difficile colitis

- Massive watery diarrhea
- On endoscopy (rarely done):
  - Pseudomembrane formation (white-yellow plaques)
  - Mucosal ulcerations, fibrin, inflammatory cells
- Diagnosis
  - Stool detection of toxin A and B

## Clostridium difficile colitis


- Treatment:
  - Metronidazole
  - ORAL Vancomycin
- Other therapy for severe, recurrent disease
  - Surgery
  - Stool transplant

## Corynebacterium diphtheria

- Causes diphtheria
- Several special features of bacteria
  - Exotoxin
  - Unique lab diagnostic techniques

## Diphtheria Exotoxin

- Not part of bacterial genome
- Carried by  $\beta$ -prophage
- "Lysogenic" phage → incorporates DNA into bacteria
- Inactivates elongation factor (EF-2)
- EF-2 necessary for protein synthesis (translation)




## Corynebacterium Features

- Gram positive rods
  - Curved
  - "Chinese character" distribution
- Special culture media required
  - Loeffler's or Tinsdale (Tellurite plate)
  - Black colonies on Tinsdale media
  - Metachromatic (blue/red) granules on Loeffler's media
- Elek test for toxin detection (for diagnosis)
  - Antitoxin-impregnated filter paper under agar
  - Bacterial toxin precipitates and can be visualized



CDC/Public Domain

## ELEK Test



## Diphtheria

- Sore throat, fever, lymphadenopathy
- Gray-white membrane in pharynx
- Absorption/dissemination of toxin can cause:
  - Myocarditis (heart failure, arrhythmias, heart block)
  - CNS disease (neuropathies)
  - Renal disease (renal failure)

## Diphtheria

- Rare due to vaccination with toxoid
- Treatment (acute infection):
  - Penicillin
  - Diphtheria antitoxin (passive immunization)
  - Diphtheria toxoid (active immunization)

## Listeria

- Found in soil
- Facultative intracellular organism
- "Tumbling motility"
- Move from cell to cell to avoid extracellular response
- Polymerizes actin in cells to move ("actin rockets")

## Listeria

- Multiplies in cells with poor cell-mediated immunity
  - Neonates, HIV, organ transplant
- In adults, often from contaminated food
  - Undercooked meat, unwashed vegetables
  - Unpasteurized cheese/milk
  - Likes cold temperatures
- In neonates, transplacental or vaginal transmission

## Listeria

- Gastroenteritis
  - Diarrhea, nausea, vomiting
  - Usually self limited
- Meningitis
  - Elderly or newborns
- Infection in pregnancy
  - Bacteremia in 3<sup>rd</sup> trimester
  - Flu-like illness (fever, chills)
  - Often resolves without treatment
  - Rarely can cause fetal demise or newborn infection

## Granulomatosis Infantiseptica

- Severe in utero infection from Listeria
- Disseminated abscesses and/or granulomas
- Multiple organs: liver, spleen, lungs, kidneys, brain
- Skin lesions (papules, ulcers)
- Most babies stillborn or die soon after birth
- Placenta shows distinctive inflammation
  - Chorioamnionitis
  - Villitis
  - Abscess formation

## Bacillus

- Bacillus anthracis
- Anthrax: Skin or pulmonary disease
  - Largely eradicated
  - Weapon of bioterrorism
- Bacillus cereus
  - Food poisoning

## Bacillus Anthracis

- Only bacteria with a polypeptide capsule
  - Most are polysaccharide
  - B. Anthracis capsule contains D-glutamate
  - Limits/prevents phagocytosis
- Found in soil
- Infects cattle, sheep, horses (and humans)
  - Farm workers at risk
- Spores can be used as bioterrorism weapon
- Produces two toxins:
  - Edema toxin
  - Lethal toxin

## Anthrax Toxins

- Edema toxin (contains edema factor)
  - Activates adenylate cyclase
  - Increases cAMP → fluid secretion
- Lethal toxin (contains lethal factor)
  - Protease
  - Inhibits cell signaling
  - Causes apoptosis

## Anthrax

- Cutaneous disease
  - Spores enter skin through cuts/abrasions
  - Vegetate → bacteria grows
  - Painless black ulcers forms
  - Can progress to bacteremia and death
  - Edema surrounds black ulcer from edema factor
- Treatment: Ciprofloxacin, Doxycycline, Clindamycin



Image courtesy of Wikipedia/Public Domain

## Anthrax

- Pulmonary disease
  - "Woolsorters' disease"
  - Inhalation of spores
  - Flu symptoms that rapidly progress
  - Pulmonary hemorrhage, mediastinitis, shock, death
- Treatment: Multi-drug regimen, antitoxin



Image courtesy of Wikipedia/Public Domain

## Bacillus Cereus

- Food poisoning from enterotoxins
- Classically in undercooked/reheated rice
  - Bacteria frequently present in uncooked rice
  - Heat-resistant spores may survive cooking
  - Cooked rice at room temperature allow bacteria to multiply
  - "Reheated rice syndrome"

## Bacillus Cereus

- Emetic type
  - Direct ingestion of toxin: Cereulide
  - Abdominal cramps, nausea, and vomiting (rarely diarrhea)
  - 1 to 5 hours after ingestion
  - Classically occurs in rice dishes
- Diarrheal type
  - Abdominal cramps and diarrhea (not vomiting)
  - 8 to 16 hours after ingestion
  - Caused by at several enterotoxins
  - Toxins are heat labile
  - Cooking food reduces risk of illness
  - Often from meats, vegetables, and sauces

## Actinomyces

- Normal oral flora
- Also found in female genital tract
- Anaerobe
- Clusters into long filaments resembling fungi
- Causes head/neck abscesses
- In women, can cause IUD infections
- Classically preceded by dental work/orofacial trauma
  - Facial mass present on exam
  - Often yellow center (yellow "sulfur" granules)
  - Often draining puss
- Treatment: Penicillin +/- drainage

## Nocardia

- Branching, filamentous (like Actinomyces)
- Acid fast
- Produces urease (can be used to identify bacteria)
- Obligate aerobe (loveslungs!)
- Found in soil


## Nocardia

- Pneumonia
  - Immunocompromised patients
  - Inhalation of bacteria
  - Many radiology findings: nodules, masses, infiltrates, lobar consolidation, pleural effusions
- Skin infection
  - Immunocompetent patients
  - Often invades skin during gardening or farming
  - Lots of manifestations: ulcers, cellulitis, nodules, abscesses
- Rarely other infections: brain abscess, bacteremia
- Drug of choice: TMP-SMX

# Gram Negative Rods

Jason Ryan, MD, MPH

## Gram Negative Bacteria



## Gram Negative Rods

- Most are in the family: enterobacteriaceae
- Many are inhabitants of the normal GI flora
- Often cause diarrhea and UTIs
- Resistant to Penicillin and Vancomycin
- Outer membrane inhibits entry of drug

## Klebsiella

- Intestinal flora
- Non-motile, capsular
- Infection with impaired host defenses
  - Alcoholics, diabetics, sick people (nosocomial)
- Infection with aspiration of GI contents
  - Aspiration pneumonia, Lung abscesses
- Often resistant to many antibiotics
- Treatment based on susceptibility testing

## Klebsiella

- Klebsiella pneumonia
  - Lobar
  - Occurs in alcoholics or diabetics, often after aspiration
  - Classically results in red "currant jelly" sputum
- Lung abscess
  - Usually caused by mouth anaerobes
  - Peptostreptococcus, Fusobacterium, Prevotella, Bacteroides
  - Can also be due to Klebsiella
- Rare cause of UTIs (3-4%)
- Liver abscesses
  - Usually in patients with underlying liver disease or cholangitis

## E. Coli

Special virulence factors

- Fimbriae (pili)
  - Attach to epithelial surfaces
  - May be specialized for surfaces (i.e. urinary tract)
- K capsule
  - K1 capsular antigen present in 75% meningitis cases (babies)
  - Inhibits phagocytosis, complement




Image courtesy of Manu Forero/Wikipedia

## E. Coli Diseases

- Watery diarrhea
- Bloody diarrhea (dysentery)
- UTI/pyelonephritis
  - E. Coli bacteremia/sepsis (rare), usually from UTI
- Meningitis in newborns

## E. Coli Diarrheal Illnesses

- 4 different E. Coli diarrheal illnesses
- EnteroINVASIVE E. Coli (EIEC)
- EnteroTOXIGENIC E. Coli (ETEC)
- EnteroPATHOGENIC (EPEC)
- EnteroHEMORRHAGIC (EHEC)

## E. Coli

- EnteroINVASIVE E. Coli (EIEC)
  - Invades intestinal mucosa
  - Necrosis, inflammation, bloody diarrhea
  - Clinically similar to Shigella (no toxins)
- EnteroTOXIGENIC E. Coli (ETEC)
  - Two toxins: Labile and Stable
  - Watery (traveler's) diarrhea (contaminated food/water)
  - No inflammation/invasion
- Enteropathogenic (EPEC)
  - No toxin, no inflammation
  - Blunt villi, prevent absorption
  - Diarrhea usually in children (p=pediatrics)

## Enterohemorrhagic E. Coli

EHEC

- Does not ferment sorbitol (sorbitol-MacConkey agar)
- Classic serotype: E. coli O157:H7
- Does not invade host cells (toxin causes disease)
- Produces Shiga-like toxin → bloody diarrhea
  - Bacteriophage-encoded (lysogenic) toxin
- Usually from undercooked beef
- Toxin Effects
  - Endothelium swells → vessel lumens narrow
  - Deposition of fibrin/platelets in microvasculature
  - Hemolysis, inflammation

## Hemolytic Uremic Syndrome

HUS

- Complicates ~10% EHEC cases
- Common in children
- Triad:
  - Hemolytic anemia
  - Thrombocytopenia
  - Acute renal failure (uremia)
- HUS + fever, mental status changes = TTP
  - Thrombotic thrombocytopenic purpura
- Usually occurs 5-7 days after diarrhea

## E. Coli Treatment

- Most E. Coli diarrheas self-limited
  - Usual treatment is hydration
  - Antiperistaltic agents (Loperamide) not helpful
  - Antibiotics rarely used (may increase toxin release)

## Gram Negative Sepsis

- Fever, tachycardia, hypotension
- Life-threatening
- Driven by endotoxin (LPS; Lipid A)
- Common scenario:
  - Elderly patient
  - UTI (catheter, BPH)
  - Gram negative sepsis (+ blood cultures)

## Infectious Diarrhea

- | Bloody   | Watery  |
|--|---|
| <ul style="list-style-type: none"><li>• Campylobacter</li><li>• Salmonella enterica</li><li>• Shigella</li><li>• Yersinia enterocolitica</li><li>• EIEC</li><li>• Entamoeba histolytica</li><li>• EHEC</li></ul> | <ul style="list-style-type: none"><li>• ETEC</li><li>• Cholera</li><li>• C. difficile</li><li>• C. perfringens</li><li>• Giardia, Crypto</li><li>• Rotavirus, Norovirus</li></ul> |

Fecal leukocytes, RBCs usually indicate invasive infection  
Mucous, epithelial cells only seen in toxin-mediated disease  
Stool ova and parasites seen in protozoal infections

## Enterobacter

- Rare cause of nosocomial UTIs
- Resistant to many antibiotics
  - Extended-spectrum beta-lactamases (ESBL)
  - Resistance to most beta-lactams: penicillins, cephalosporins, and aztreonam
- Often treated with Carbapenems
  - Imipenem, Meropenem

## Citrobacter & Serratia

- Slow lactose fermenters
- Not dominant pathogen for any clinical condition
- Often resistant to many antibiotics
- Citrobacter
  - Can be found in normal GI flora
  - Gram negative sepsis (with other GN bugs)
- Serratia
  - Produce distinctive red colonies (red pigment)
  - Catalase positive
  - Hospital outbreaks: contaminated water, soap, IV solutions
  - Sometimes osteomyelitis in IV drug users (also pseudomonas)

## Salmonella

- Two general types
  - Salmonella typhi → typhoid fever
  - Non-typhoid strains → Enterica, enteritidis
- Non typhoid strains cause invasive gastroenteritis
  - Nausea, vomiting, cramps, bloody diarrhea
  - Ingestion of contaminated meat, eggs, poultry

## Salmonella

- Flagellated and motile
- Encapsulated
- Disseminate through blood
  - Osteomyelitis in sickle cell patients
- Live in GI tract of mammals, birds, reptiles
- Produce hydrogen sulfide
  - Triple Sugar Iron (TSI) test → media turns black
  - Differentiates from Shigella
- Invades GI mucosa
- Cellular response: Largely monocytes

## Typhoid Fever

Salmonella Typhi

- Fever, headache, abdominal pain, diarrhea
- Travelers to Asia, Africa, South America
- Classic feature #1: Rose spots
  - Faint salmon-colored macules
  - Trunk and abdomen
- Classic feature #2: Pulse-temperature dissociation
  - High fever → slow pulse
- Can remain in gall bladder (carrier state)
  - Endemic countries 1-4% people may be carriers
  - May be risk factor for carcinoma

## Salmonella

- Diagnosis: Culture (stool, blood)
- Treatment gastroenteritis:
  - Fluids/electrolytes
  - Few data showing antibiotics are helpful (may prolong illness)
  - Difficult to treat: Lots of antibiotic resistance
  - Antibiotics used in severely ill patients only
  - Anti-peristalsis meds (Loperamide) contraindicated
- Typhoid fever: Ceftriaxone, Fluoroquinolones
- Typhoid vaccine available
  - Inactive variant of bacteria given orally
  - Used for traveler's to high risk areas

## Shigella

- Nonmotile (no flagella)
- Invades mucosal cells (M cells in Peyer's patches)
  - Macropinocytosis
  - Induces apoptosis
- Spreads from cell to cell
  - Does not spread via bloodstream (like Salmonella)
- Releases Shiga toxin
  - But non-toxin strains still cause disease
  - Cellular invasion more important mechanisms of disease
- Very few bacteria can cause disease (few as 10!)
- Cellular response: Largely PMNs

## Shigella

- Not normal GI flora
- Fecal-oral transmission
- Common in children
- Diagnosis: Stool Culture
- Treatment:
  - Fluids/electrolytes
  - Antibiotics improve symptoms, reduce shedding in stool
    - Can limit spread
  - Ceftriaxone or Ciprofloxacin

## Salmonella and Shigella

Both GNRs, both cause bloody diarrhea, both invasive

Salmonella	Shigella
H <sub>2</sub> S (black on TSI)	No H <sub>2</sub> S
Monocytes	PMNs
Hematogenous spread	Cell to cell spread (Macropinocytosis)
No antibiotics	Yes antibiotics
Motile (flagella)	Nonmotile
--	Shiga toxin
--	Low infectious dose

## Proteus

- Rare cause of UTIs
- "Swarm phenotype"
  - Long flagella
  - Facilitates urinary ascent
  - Bulls-eye on agar plates
- Produces urease
  - Converts urea to ammonia
- Associated with struvite kidney stones

## Pseudomonas aeruginosa


- Highly feared bacteria
  - Serious hospital acquired infections (i.e. VAP)
  - Resistant to many antibiotics (“anti-pseudomonal drugs”)
- Many infections:
  - Pneumonia (nosocomial)
  - UTIs
  - Surgical wound infections
  - Bacteremia/sepsis
  - Osteomyelitis
  - Otitis externa

## Pseudomonas aeruginosa

- Obligate aerobe
- Loves the water
  - Commonly found in environmental water sources
- Produces a blue-green pigment (pyocyanin)
  - Smooth, florescent green colonies on culture media
- Sweet, grape-like odor

## Pseudomonas Toxins

- Endotoxin (LPS)
  - Can cause fever, shock
- Exotoxin A
  - Inactivates elongation factor (EF-2)
  - EF-2 necessary for protein synthesis



## Pseudomonas Infections

- Pneumonia in cystic fibrosis
- Burn infections
- Hot tub folliculitis
  - Tender, itchy papules after using a hot tub or spa
- Otitis externa (Swimmer’s ear)
  - Inflammation of the outer ear and ear canal
  - Painful ear with discharge

## Pseudomonas Infections

- Osteomyelitis in IV drug users
- Ecthyma gangrenosum
  - Black, necrotic ulcers on skin with bacteremia
  - Invasion/destruction blood vessels by bacteria
  - Classic case is neutropenic cancer patient with fever, chills (bacteremia) who develops black lesions on chest/back

## Burkholderia cepacia

- Gram negative rod similar to pseudomonas
- Oxidase positive
- Catalase positive
- Rare cause of infections:
  - Cystic fibrosis
  - Chronic granulomatous disease

## H. Pylori

- Causes gastritis and ulcers (abdominal pain)
- Recently identified bacteria (1982!)
- Urease positive
  - Hydrolyzes urea to compounds that damage epithelium
  - Produces ammonium (alkaline)
  - Protects bacteria from stomach acid
- Urea breath test
  - Patients swallow urea with isotopes (carbon-14 or carbon-13)
  - Detection of isotope-labelled carbon dioxide in exhaled breath
  - Indicates urea was split (i.e. urease present)

## H. Pylori

- Infection common in patients with ulcers
  - Majority of patients with duodenal ulcers
  - Many patients with gastric ulcers
- MALT lymphoma
  - Mucosal associated lymphoid tissue
  - B-cell cancer, usually in the stomach
  - HIGHLY associated with H. Pylori infection
- Diagnosis:
  - Biopsy
  - Urea breath test
  - Stool antigen

## H. Pylori

- Treatment: "Triple therapy" for 7-10 days
  - Proton pump inhibitor
  - Clarithromycin
  - Amoxicillin/Metronidazole
- Testing often repeated to confirm eradication
  - Breath test, stool antigen, or biopsy
- Treatment failures ~20%
  - Alternate regimens can be tried

## Legionella

- Does not gram stain well
- Silver stains used
- Special culture requirements
- Buffered charcoal yeast extract agar (BCYE)
- Iron and cysteine added for growth
- Supplemented with antibiotics and silver dyes
  - Antimicrobials prevent overgrowth by competing organisms
  - Dyes give distinctive color to Legionella

## Legionella

- First identified at American Legion convention
- Infection from inhalation of aerosolized bacteria
  - Not airborne
- Outbreaks at hotels with contaminated water
- Can cause nosocomial pneumonia in nursing homes

## Legionella

### Symptoms

- Initially mild pneumonia symptoms
  - Fever; mild, slightly productive cough
- Can progress to severe pneumonia
- GI symptoms
  - Watery diarrhea, nausea, vomiting, and abdominal pain
- Hyponatremia (Na<130 meq/L) common
  - Can occur in any PNA but more common Legionella

## Legionella

### Diagnosis

- Classic Case
  - Mild cough
  - Watery diarrhea
  - Confusion (low Na)
  - Negative bacteria on gram stain
- Treatment: Fluoroquinolone or Macrolide

## Pontiac Fever

- Mild form of Legionella infection
- Fever, malaise, chills, fatigue, and headache
- No respiratory complaints
- Chest radiograph usually normal

## Bacteroides fragilis


- Gram (-) rod
- Anaerobic bacteria
- Normal GI flora
- Cause infection after breach of mucosal barrier

## Bacteroides fragilis

- Rarely causes infections alone
- Usually part of polymicrobial infections from GI tract
  - Peritonitis (following perforation)
  - Intraabdominal abscess
  - Lung abscess (aspiration)
  - E. Coli/GNRs and B. Fragilis often components together
- Covered by metronidazole
- Common GI therapy: Cipro/Flagyl
  - Quinolone for E. Coli
  - Metronidazole for B. Fragilis

## Other Gram Negatives

Jason Ryan, MD, MPH



## Moraxella catarrhalis

- Gram negative diplococci
- Colonizes airway
- Can cause otitis media, COPD exacerbations
- Usually treated empirically without micro diagnosis
- Not in most micro lab algorithms
  - Most labs will not speciate airway samples with gram (-) cocci
  - Non-virulent strains Neisseria normal airway flora

## Neisseria

- Meningitis and Gonorrhea
- Both gram negative cocci in pairs (diplococci)
- Both ferment glucose
- Meningococcus ferments Maltose
- Gonococcus ferments only Glucose
- Both produce IgA protease
- Ceftriaxone often used to treat both

## Neisseria Meningitidis

- Causes meningitis and meningococemia
- Transmitted by respiratory droplets
- Enters pharynx then bloodstream then CSF
- Many asymptomatic carriers

## Neisseria Meningitidis

- Polysaccharide capsule prevents phagocytosis
- Lipooligosaccharide (LOS) outer membrane
  - Like LPS on enteric gram negative rods
  - Endotoxin → many toxic effects on body
  - Activates severe inflammatory response
- Vaccine available
  - Contains capsular polysaccharides → anti-capsule antibodies

## Neisseria Meningitidis


- Bacteremia can complicate meningitis
  - Meningococemia
- Sepsis: fevers, chills, tachycardia
- Purpuric rash
- DIC
- Waterhouse-Friderichsen syndrome
  - Adrenal destruction from meningococemia
- Life-threatening

## Neisseria Meningitidis

- Can cause outbreaks
  - Dorms, barracks
- Can infect young, healthy people
  - College students in dorms
- Infected patients need droplet precautions
- Close contacts receive prophylaxis/vaccine
  - Rifampin
  - Also Ceftriaxone or Ciprofloxacin

## Neisseria Meningitidis

Terminal complement pathway deficiency



C5-C9 Deficiency  
Recurrent NM Infections  
Most often meningitis

Wikipedia/Public Domain

## Neisseria Gonorrhoea

- Mainly causes gonorrhoea (STD)
- Can also cause:
  - Pelvic inflammatory disease (PID)
  - Septic arthritis
  - Neonatal conjunctivitis
  - Fitz-Hugh-Curtis syndrome
- Key feature: antigenic variation of **pilus proteins**
  - No long term immunity after infection
  - Re-infection likely
  - No vaccine

## Gonorrhoea

- Most men/women with N. Gonorrhoea asymptomatic
- N. Gonorrhoea and Chlamydia often co-infect
  - Both can cause same symptoms
  - Treat for both (Ceftriaxone, Azithromycin/Doxycycline)
- Men: Urethritis
  - Dysuria, discharge
  - Can progress to or epididymitis/orchitis
- Women: Cervicitis
  - Itching, discharge from cervix
  - Not painful
  - Can progress to PID

## Gonorrhoea

- Pelvic inflammatory disease
  - Infection ascends (uterus, ducts, ovaries)
  - Pelvic/abdominal pain
  - Dyspareunia
  - Cervical motion tenderness on exam (chandelier sign)
  - High risk of subsequent **ectopic pregnancy, infertility**
- Fitz-Hugh-Curtis
  - Perihepatitis
  - Inflammation of Glisson's capsule around liver
  - Severe RUQ tenderness with pleuritic pain
  - "Violin string" adhesions of parietal peritoneum to liver

## Septic Arthritis

- Disseminated gonococcal infection (0.5 to 3%)
- Septic arthritis
- Key scenario:
  - Sexually active young person
  - Swollen, warm and painful knee

## Neonatal Conjunctivitis

- Ophthalmia neonatorum
- Can also be caused by Chlamydia
- Swelling and discharge from eye
- 5 to 14 days after birth
- Untreated can lead to visual impairment
- Prophylaxis: Erythromycin ophthalmic ointment
- Newborn prophylaxis mandated by many states

## Chlamydia

- Obligate intracellular organisms
  - Cannot make their own ATP
- Cell wall lacks muramic acid
  - *N*-acetylmuramic acid (NAM) in peptidoglycan
  - Cell wall lacks peptidoglycan
- Do not gram stain well (technically gram negative)
- Giemsa stain

## Chlamydia

- Penicillins do not work well
- Ceftriaxone (for Gonorrhea) ineffective
- Treatment of choice: azithromycin, doxycycline
  - Protein synthesis inhibitors

## Chlamydia

- Two phases to life cycle
- #1: Elementary body (small, dense)
  - Enters cell via endocytosis
- #2: Reticulate body
  - Replicates in cells by fission
  - Can be seen in tissue culture
- Elementary bodies and reticular bodies grow, multiply, eventually rupture cell and disperse

## Chlamydia

- Chlamydia trachomatis (sexually transmitted)
  - Nongonococcal urethritis
  - PID
  - Conjunctivitis
  - Reactive arthritis
- Chlamydochlamydia pneumoniae
  - Atypical pneumonia
  - Transmitted by aerosol
- Chlamydochlamydia psittaci
  - Psittacosis (Parrot fever)
  - Infection from birds

## Chlamydia trachomatis

- Sexually transmitted
- Often asymptomatic in men & women
- Women: Cervicitis
  - Discharge, post-coital bleeding
  - Can progress to PID, Fitz-Hugh-Curtiss
- Men:
  - Discharge, dysuria
- Treatment: Azithromycin/Doxycycline
  - Plus Ceftriaxone for N. Gonorrhea

## Chlamydia trachomatis

### Newborns

- Infection from passage through birth canal
- Conjunctivitis
  - Similar to Gonorrhea
- Pneumonia
  - 4-12 weeks old
  - Classic feature is "staccato cough"
  - Inspiration between each single cough
  - Often have a history of conjunctivitis

## C. Trachomatis Diagnosis

- Nucleic acid amplification testing (NAAT)
  - PCR of Chlamydia DNA/RNA
  - Gold standard
- Culture and staining
  - No longer done routinely
  - Giemsa stain
  - Chlamydial inclusion bodies in cytoplasm of epithelial cells

## Reactive Arthritis

- Autoimmune arthritis
- Triggered by infection
- Intestinal infections
  - Salmonella, Shigella, Campylobacter, Yersinia, C. Difficile
- Chlamydia trachomatis
- Classic triad (Reiter's syndrome)
  - Arthritis (often unilateral, lower extremities, knees, toes)
  - Conjunctivitis (red eye, discharge)
  - Urethritis (dysuria, frequency - noninfectious)
- Diagnosis: Classic features following typical infection
- Treatment: NSAIDs

## Lymphogranuloma Venereum

- Chlamydia infection that enters lymphatics
- Different serotypes from those that cause urethritis
- Sexually transmitted
- Initially: Genital ulcer
  - Sometimes unnoticed; Resolves
- Later: Tender inguinal or femoral lymph nodes
- Treatment:
  - Drainage
  - Antibiotics



Herbert L. Fred, MD and Hendrik A. van Dijk

## Trachoma

### Chlamydia Eye Disease

- Caused by unique serotypes of C. Trachomatis
- Mostly in Africa and other developing parts of world
- Highly contagious
- Spread by contact with eye secretions
- Acutely causes conjunctivitis
- Repeated infections → Corneal scarring → blindness
- Leading cause of infectious blindness worldwide

## C. Trachomatis Serotypes

Serotype	Infections
A, B, C	Trachoma, Blindness; Found in Africa
D through K	Urethritis, PID, neonatal pneumonia, neonatal conjunctivitis
L1, L2, L3	Lymphogranuloma venereum

## Campylobacter



Wikipedia/public domain

- Usually *C. jejuni*, sometimes *C. coli*
- Faint, gram negative bacteria
  - Gram stain not sensitive
- Spiral shaped, curved rod (comma shaped)
- Oxidase positive
- Microaerophilic
  - Requires oxygen but lives best with low O<sub>2</sub> levels

## Campylobacter

- VERY common cause acute diarrhea in children
- A leading cause of acute diarrhea worldwide
- Fecal-oral transmission
  - Lives in animal intestines, especially poultry
  - Undercooked meat especially poultry
  - Unpasteurized milk
  - Can also contaminate drinking water
- Common trigger of Guillain-Barre
  - Demyelinating disease
  - Ascending weakness

## Vibrio Cholera

- Toxin-mediated disease
- Toxin carried by bacteriophage (lysogenic)
- Permanently activates G<sub>s</sub> → ↑cAMP
- Voluminous "rice-water" diarrhea

## Vibrio Cholera

- Requires large "dose" of bacteria for infection
  - Acid kills small amounts of bacteria
  - Common in areas that lack clean water
  - Also can occur on acid suppression drugs (lowers dose req'd)
- Death from dehydration, electrolyte losses, shock
- Treatment: Aggressive volume repletion

## V. vulnificus/parahaemolyticus

- Both cause food poisoning (diarrhea)
- Found in raw oysters
- *V. vulnificus* can infect wounds after swimming in contaminated water

## Haemophilus influenzae

- Colonizes nasopharynx
- Causes several respiratory diseases, meningitis
- Some have polysaccharide capsule some do not
  - "Capsular" bacteria are "typeable" into six serotypes (a to f)
  - Others are "nontypeable"
- Most disease caused by type B
  - Capsule consists of ribosyl and ribitol phosphate polymer
- Vaccine contains type B capsule (Hib)
  - Conjugated to a carrier protein (often tetanus toxoid)
  - Stronger T-cell response
  - Given before 7 months

## Haemophilus influenzae

### Special Features

- IgA protease
- Grows on Chocolate agar
  - Factors V (NAD) and X (hematin) present
  - Will also grow with S. Aureus on blood agar

## Haemophilus influenzae

### Infections

- Epiglottitis
  - Life-threatening (airway obstruction)
  - Unvaccinated children with fever, sore throat
  - Dysphagia, drooling
  - Epiglottitis will appear "cherry red"
- Pneumonia
- Meningitis
- Otitis media, bronchitis, conjunctivitis
  - S. pneumoniae and non-typeable H. influenzae
  - Vaccine not protective

## Haemophilus ducreyi

- Causes chancroid
- Painful genital ulcer
- Contrast with chancre (syphilis): non-painful
- Sexually transmitted
- Treatment: Azithromycin/Ceftriaxone

## Genital Ulcers

Disease	Feature
Syphilis	Painless chancre
H. Ducreyi	Painful chancroid
Herpes	Multiple vesicles/ulcers
Lymphogranuloma Venereum	Large, swollen lymph nodes (buboes)

## Bordetella Pertussis

- Causes whooping cough
- URI with severe coughing
- Classic presentations
  - Paroxysms of coughing
  - Inspiratory "whoop"
  - Post-cough vomiting
  - Exhaustion from coughing
- Coughing fits can last weeks
- In China, pertussis known as the "100 day cough"

## Bordetella Pertussis

- Transmitted by aerosolized droplets
- Pertussis toxin
  - Shown to inhibit G<sub>i</sub> proteins
  - Allows over-activation of adenylate cyclase
  - ↑cAMP levels in cells in neutrophils
  - Result: impaired recruitment of neutrophils
- Toxin may not be cause of cough
  - Some species without toxin shown to cause symptoms

## Bordetella Pertussis

- Infection rare due to vaccine
- Acellular pertussis vaccines used
- Contain purified pertussis antigens

## Yersinia Enterocolitica

- Found in domesticated animals (dogs), pigs
- Often transmitted through contaminated pork
- Also from contaminated water or milk
- Fever, abdominal pain, nausea, vomiting
- Bloody diarrhea
- Can cause inflammation around appendix or in mesenteric lymph nodes (mesenteric adenitis)
  - May mimic Crohn's or appendicitis
- Don't confuse with Yersinia pestis (plague)

# Spirochetes

Jason Ryan, MD, MPH

## Spirochetes

- Bacteria with long, corkscrew-shaped cells
- Difficult to grow/culture
- Serology usually used for diagnosis




Image courtesy Wikipedia/Public Domain

## Spirochetes

- Leptospira (Leptospirosis)
- Borrelia (Lyme disease)
- Treponema (Syphilis)

## Leptospira interrogans

- Causes Leptospirosis
- Lives in rodents → shed in urine
- Illness commonly from contaminated water
- Disease ranges asymptomatic → severe
- Flu-like illness: fever, rigors, myalgias, headache
- Conjunctival suffusion (red eyes)
- Aseptic meningitis
- Weil's disease (rare complication)
  - Liver damage (jaundice), renal failure, and bleeding

## Leptospira interrogans

- **Diagnosis:**
  - Usually from classic history/exam
  - Serology (antibody) tests available
- **Treatment: Doxycycline or Azithromycin**
- **Classic case:**
  - Surfer or swimmer in Tropics
  - Flu-like illness
  - Conjunctival suffusion
  - Jaundice

## Borrelia burgdorferi

- Causes Lyme disease
- Cause by tick bite (*Ixodes scapularis*)
  - Tick larvae feed on mice (reservoir for *Borrelia*)
  - Infected adult ticks feed on deer
- Ticks can bite humans → infection with *Borrelia*
- Tick must be attached ~48hrs to transmit bacteria
- Common in Northeast US (Lyme, Connecticut)

## Lyme Disease

- Stage 1: Erythema chronicum migrans
  - Classic finding: expanding "Bull's-eye" rash
  - Flu-like symptoms
- Stage 2: Neurologic and cardiac
  - Facial nerve palsy
  - AV block



Image courtesy of Wikipedia/Public Domain



## Lyme Disease

- Stage 3:
  - Arthritis (often knees)
  - Neuropathy (pain, paresthesias)
  - Encephalopathy (mild cognitive disturbance)
  - Rash: Blue-red discoloration
    - Acrodermatitis Chronica Atrophicans
    - More common in European Lyme
- Treatment: Doxycycline or Ceftriaxone

## Relapsing Fever

- US: *Borrelia hermsii* and *Borrelia turicatae*
  - Transmitted by tick bites
- Developing world: *Borrelia recurrentis*
  - Transmitted by louse (insect)
  - Spread from person to person by louse (epidemics)
- Symptoms are relapsing fever (duh!)
- Antigenic variation causes recurrent fevers
- Spirochetes change major antigens on surface
- This evades immune response
  - Growth occurs
  - Fever returns

## Treponema pallidum

- Causes syphilis
- Sexually transmitted disease
- Can see spirochete by dark field microscopy
- Disease: 1<sup>o</sup>, 2<sup>o</sup>, 3<sup>o</sup> stages




Image courtesy of Wikipedia/Public Domain

## Primary Syphilis

- Two to three weeks after exposure
- Painless chancre (ulcer)
  - Classically on the penis
  - Usually 1-2cm
  - Raised
- Often unnoticed (painless) → disease progresses



Image courtesy of Wikipedia/Public Domain

## Secondary Syphilis

- Rash
  - Classically maculopapular rash
  - Covers all extremities including palms/soles
- Flu-like symptoms
  - Fever, headache, malaise, sore throat, myalgias
- Condyloma lata
  - Large, raised, gray to white lesions
  - Moist areas: inside mouth, perineum
  - Often close to chancre; may reflect direct spread
- Treponema present in condyloma and chancre
  - Can visualize with dark-field microscopy

## Palms and Soles


- Most maculopapular rashes spare palms/soles
- Three that don't:
  - Syphilis
  - Rock Mountain Spotted Fever
  - Coxsackie virus (hand, foot, mouth)

## Tertiary Syphilis

- Gummas
  - Form of granuloma
  - Mass lesions that can appear anywhere: skin, liver
  - Often mistaken for tumors
- Aortitis
  - Vasa vasorum inflammation
  - Risk of aortic dissection
- Neurosyphilis
  - Many, many symptoms
  - Meningitis, dementia, nerve palsies

## Tertiary Syphilis

- Argyll Robertson pupil
  - "Prostitute's pupil"
  - Small pupils
  - Constrict to accommodation
  - Do not constrict to light
- Tabes Dorsalis
  - Demyelination of posterior columns
  - Wide-based gait
  - Ataxia (falls, loss of balance)



## Syphilis Diagnosis

- VDRL
  - Venereal Disease Research Laboratory
  - "Non treponemal" test
  - Serum reacted with cardiolipin antigen (nonspecific)
- RPR
  - Rapid Plasma Reagin
- Syphilis patients' serum (antibodies) will react
  - Positive VDRL/RPR
- For neurosyphilis need to test CSF
  - Positive serum test does not necessarily indicate CNS disease
  - Many false negatives – difficult diagnosis

## Syphilis Diagnosis

- Many false positives VDRL/RPR
  - Mononucleosis
  - Rheumatic fever
  - SLE
  - Leprosy
  - Pregnancy

## Syphilis Diagnosis

- FTA-ABS
  - Fluorescent treponemal antibody absorption
  - "Treponemal test"
  - Detects antibodies against specific treponemal antigens
  - Very specific
- Test result: "Reactive" "Non-reactive"

## Congenital Syphilis

- Spirochete transmitted from mother to fetus
- Can occur in 1<sup>st</sup> trimester
  - Mothers screened early in pregnancy
- Most often in mothers with no prenatal care
- Findings on baby can be early or late
  - Early (<2yrs); Late (>2yrs)

## Congenital Syphilis

### Early Findings

- Hepatomegaly
- Runny nose
- Maculopapular rash
  - Small, red or pink spots
  - Often on back, buttocks, posterior thighs, and soles
- Abnormal long-bone radiographs
  - Many, many abnormalities reported

## Congenital Syphilis

### Late Findings

- Ears/nose
  - Saddle nose (no nasal bridge)
  - Hearing loss/deafness
- Teeth
  - Hutchinson teeth (notched, peg-shaped teeth)
  - Mulberry molars (maldevelopment of the molars)
- Legs
  - Saber shins (bowed legs)

## Syphilis Treatment

- Penicillin G
- Jarisch-Herxheimer reaction
  - Flu-like syndrome after starting antibiotics
  - Killed bacteria cause immune response
  - Self-limited

# Zoonotic Bacteria

Jason Ryan, MD, MPH

## Zoonotic Bacteria

- All rare, all transmitted from animals to humans
- Almost all can be treated with Doxycycline
- Key is to recognize clinical syndromes

<u>Animal Vector</u>	<u>Tick-Flea-Louse</u>
Bartonella	Ehrlichia
Brucella	Borrelia
Chlamydia psittaci	Rickettsia Rickettsia
Coxiella burnetii	Rickettsia Typhus
Leptospira	Rickettsia Prowazekii
Francisella tularensis	Yersinia pestis
Pasteurella	

## Bartonella henselae

- Cats harbor fleas that carry Bartonella
- Cat Scratch Fever
  - Cat scratch (almost always a child)
  - Red, swollen area 3-10 days later
- Regional lymphadenopathy (hallmark of disease)
  - Tender, red lymph nodes

## Bartonella henselae

- Bacillary Angiomatosis (vascular lesions)
  - Diffuse skin papules in AIDS patients
  - Often mistaken for Kaposi Sarcoma
- Endocarditis (rare cause, culture negative)
- Treatment: Doxycycline, Azithromycin

## Kaposi Sarcoma

- Raised, red/purple skin lesions
- Common in HIV/AIDS
- Angioproliferation
- Caused by HHV-8
- Can have similar appearance to Bacillary Angiomatosis
- Key differences
  - Kaposi Sarcoma: Lymphocytes
  - BA: Neutrophils/lymphocytes



## Granulomatous Infections

- Tuberculosis
- Leprosy
- Fungal pneumonias (Histo, Blast, Coccidio)
- Bartonella (cat scratch disease)
- Brucella
- Listeria in infants (Granulomatosis Infantiseptica)
- Schistosomiasis (worm)
- Syphilis (gummas)

## Brucella

- Lives in cows, goats
- Infection from unpasteurized milk or animal exposure
- Classic patients:
  - Worker in meat packing plant
  - Traveler from Mexico who consumed milk/cheese
- Brucellosis (undulant fever)
  - Flu-like illness
  - High fever that rises and falls
  - Profuse sweating
- Treatment: Doxycycline + streptomycin/rifampin

## Chlamydomphila psittaci

### Psittacosis

- Parrot fever
- Infection from inhalation dried feces
- Classic patient: Pet store employee
- Fever, headache, and dry cough
- Treatment: Doxycycline

## Coxiella burnetii

### Q fever

- Farm animals: cattle, goats, sheep
- Forms spores that get inhaled
- High concentrations in placenta of infected animals
- Symptoms
  - Pneumonia with flu symptoms (fever, headache, myalgias)
  - Endocarditis
- Diagnosis: Serology (antibodies)
- Treatment: Doxycycline

## Culture Negative Endocarditis

- Evidence of endocarditis with sterile BCx
- Coxiella burnetii
  - Q fever
  - Farm animals (cattle, sheep, goats)
- Bartonella
  - Cat scratch fever
  - Cat fleas

## Francisella tularensis

### Tularemia (Rabbit fever)

- Important reservoirs: Ticks, deer flies, rabbits
- Occurs in animal handlers, especially rabbits
  - Also from tick bites
- Ulceroglandular tularemia (most common form)
  - Fever, chills malaise
  - Classically the fever abates for few days, returns
  - Skin ulcer at site of insect bite
  - Swollen, painful lymph nodes
- Treatment: Streptomycin (Doxycycline okay, too)

## Pasteurella

- Lives in mouth of cats and dogs
- Infection: Cat/dog bites or scratches
- Key infections:
  - Cellulitis
  - Osteomyelitis
- Bite wounds usually polymicrobial (S. Aureus)
- Broad spectrum empiric therapy
  - Amoxicillin-clavulanate (oral)
  - Ampicillin-sulbactam (IV)
  - Piperacillin-tazobactam (IV)

## Leptospira interrogans

- Causes Leptospirosis
- Lives in rodents → shed in urine
- Illness commonly from contaminated water
  - Classic case is surfer or swimmer in tropics
- Treatment: Doxycycline or Azithromycin

## Ehrlichia

### Ehrlichiosis

- Tick-borne illness (Lone Star tick)
- White tail deer are principal reservoir
- Obligate intracellular bacteria
  - “Berry like” inclusions in monocytes (morulae)
- Symptoms
  - Flu-like illness
  - Leukopenia
  - Thrombocytopenia
- Diagnosis: Giemsa stain, serology
- Treatment: Doxycycline




Image courtesy of Wikipedia/Public Domain

## Anaplasma

### Anaplasmosis

- Bacteria very similar to Ehrlichia
- Morula seen in granulocytes (not monocytes)
- Tick vector: *Ixodes scapularis* (not Lone Star tick)
  - Same vector as Lyme disease, Babesiosis
- Similar symptoms, treatments to Ehrlichiosis
  - Fever, joint pains
  - Low WBCs, platelets
  - Blood smear: granulocytes (not monocytes) with inclusions

## Borrelia burgdorferi

- Causes Lyme disease
- Cause by tick bite (*Ixodes scapularis*)
  - Tick larvae feed on mice (reservoir for *Borrelia*)
  - Infected adult ticks feed on deer
- Ticks can bite humans → infection with *Borrelia*
- Common in Northeast US (Lyme, Connecticut)
- Treatment: Doxycycline or Ceftriaxone

## Rickettsia and Chlamydia

- Similar types of bacteria
- Obligate intracellular bacteria
  - Use host ATP
  - Chlamydia cannot make ATP
  - Rickettsia can make some
- Cannot be cultured on common media
  - Inoculated into living cells (yolk sac of chicken embryos)
- Very small
  - Close to size of viruses
- Chlamydia: Person to person
- Rickettsia: Tick- or insect-borne illnesses

## Rickettsia

- All infections occur from ticks-fleas-lice
- Three subtypes → different infections
  - *R. rickettsii* (Rocky Mountain Spotted Fever)
  - *R. typhi* (Murine typhus)
  - *R. prowazekii* (Epidemic typhus)

## Typhus versus Typhoid

- Typhus = Greek word for smoky or hazy
  - Used by Hippocrates to describe state of mind
  - Typhus caused by Rickettsia sp.
  - Can cause plagues (R. prowazekii)
- Typhoid Fever
  - Caused by Salmonella typhi
  - Enteric disease
  - Fever, diarrhea, rose spots

## Rocky Mountain Spotted Fever

Rickettsia rickettsii

- Occurs throughout US (despite name)
- Transmitted by tick bite
  - 1/3 may not recall/notice the bite
- Triad: Headache, fever, rash
- Headache, fever often come first
- Maculopapular rash
  - Starts wrists/ankles → spreads to trunk, palms, soles
- Rarely complications:
  - Encephalitis
  - Seizures
  - DIC
- Treatment: Doxycycline

## Murine Typhus

Rickettsia typhi

- Also called "endemic" typhus
  - Endemic to certain populations (no epidemics)
- Reservoir: Rats
- Transmitted from ratfleas
- Common in developing world
- Flu-like illness
- Rash (<50%)
  - Maculopapular
  - Starts in trunk spreads out
- Treatment: Doxycycline

## Epidemic Typhus

Rickettsia prowazekii

- Mostly historical
- Epidemics throughout history have killed millions
- Some outbreaks in Africa during civil wars
- Transmitted by body louse
  - Body louse lives on skin/clothes
  - Eggs laid on clothes and hatch
  - Larvae suck blood
  - During meal, louse defecates highly infective feces
  - Rickettsia in louse feces introduced to skin/membranes

## Epidemic Typhus

Rickettsia prowazekii

- Fever, chills, headaches, malaise
- Maculopapular rash
  - Starts in trunk spreads out
- Confusion, seizures, coma
- Treatment: Doxycycline

## Yersinia pestis

Bubonic Plague

- Reservoir: rats, sometimes squirrels or prairie dogs
- Humans get disease from rat flea bites
- Human to human spread via respiratory droplets
- Fever, chills, headache
- Intense pain/swelling of a lymph node area (bubo)
- Buboes:
  - Exquisite tenderness
  - Erythema and edema of overlying skin
  - Inguinal region most frequent ("bubo" = Greek word "groin")
- Treatment: Streptomycin (Doxycycline okay, too)

## Tick-Flea-Louse

- Ehrlichia → Ehrlichiosis
- Borrelia → Lyme disease
  - \*\* Babesia (parasite) → same tick
- R. Rickettsia → Rocky Mountain Spotted Fever
- R. typhi → Murine typhus
- R. prowazekii → Epidemics/plague
- Y. Pestis → Bubonic plague

# Mycobacteria

Jason Ryan, MD, MPH

## Mycobacteria

- Mycolic acids in cell wall
- Lipid-rich cell wall that is "acid fast"
- Resistant to decolorization by acid after staining with carbolfuchsin
- Do not gram stain well (technically gram positive)

## Ziehl-Neelsen

- The "acid fast" stain
- Contains carbolfuchsin
- Used to detect mycobacterium (especially TB)
- Also used for Nocardia




Image courtesy Wikipedia/Public Domain

## Mycobacteria

- M. Tuberculosis
- M. avium complex (MAC)
- M. kansasii
- Mycobacterium leprae

## MAC

Mycobacterium avium complex

- Most common non-TB mycobacterial infection
- Includes several bacteria:
  - M. avium, M. intracellulare
- Slow growing, acid-fast organisms

## MAC

Mycobacterium avium complex

- Found in water and soil
- Inhaled or ingested
- Very rare cause of pulmonary disease in non-HIV
- HIV/AIDS: Disseminated disease
  - Very low CD4 count (<50)
  - Fever, sweats, abdominal pain, diarrhea, weight loss
  - Severe anemia
  - Hepatosplenomegaly
  - ↑ alk phos, ↑ LDH
  - Often no lung findings (if lung findings → it's TB)

## MAC

Mycobacterium avium complex

- Diagnosis: Blood culture (takes 7 days or more)
- Treatment:
  - Clarithromycin plus Ethambutol
- Prophylaxis: Azithromycin

## M. Kansasii

- Most frequent non-TB mycobacteria after MAC
- Environmental source not clear
- Similar pathology but less virulent than TB
  - Fever, sweats, cough, dyspnea
  - CXR infiltrates
- Treatment: Similar to TB

## M. Leprae

Leprosy (Hansen's Disease)

- Obligate intracellular organism
- Grows very slowly - cannot be cultured
- Grows best at cool temps (27 to 33°C)
  - Infection involves skin
  - Extremities, face
- Reservoir is armadillos
- Mode of transmission unclear
- Causes granulomatous inflammation
- Mostly found in developing countries
- Most US cases occur in immigrants

## M. Leprae

Leprosy (Hansen's Disease)

- Infects skin and superficial nerves
- Key signs/symptoms
  - Skin lesions
  - Loss of sensation

## M. Leprae

Leprosy (Hansen's Disease)

- Spectrum of disease
- Severity based on strength of cell-mediated response
- Tuberculoid leprosy - Milder disease
- Lepromatous leprosy - Severe disease

## Tuberculoid Leprosy

- Patches of hypopigmented skin
- Loss of sensation over affected area
- Strong cell-mediated TH1 response contains infection
- Lesions show granulomas, few bacteria

## Lepromatous Leprosy

- Diffuse skin lesions
- Often deformed, thickened skin
- Hypopigmentation and hair loss
- Severe neuropathy (weakness, regional anesthesia)
- Th2 response
  - Humoral immunity
  - Depressed cell-mediated immunity
  - Antibodies cannot reach intracellular bacteria
- Lesions:
  - Multiple bacteria
  - No granulomas



Wikipedia/Public Domain

## M. Leprae

Leprosy (Hansen's Disease)

- **Diagnosis:**
  - Acid-fast organisms on skin biopsy
  - Note: False positive VRDL
- **Treatment:**
  - Tuberculoid: dapsone and rifampin (6 months)
  - Lepromatous: dapsone, rifampin, and clofazimine (years)

## M. Leprae

Leprosy (Hansen's Disease)

- **Rifampin**
  - Tuberculosis drug
  - Blocks RNA synthesis
- **Dapsone**
  - Competes with bacterial para-aminobenzoic acid (PABA)
  - Inhibits dihydropteroate synthetase
  - Disrupts folic acid pathways (like sulfonamides)
  - Also used for pneumocystis jiroveci (like sulfonamides)
  - Hemolysis in G6PD (like sulfonamides)
  - Rarely can cause agranulocytosis (ANC=0)
- **Clofazimine**


## IL-12 Receptor Deficiency

- IL-12 triggers differentiation T-cells to Th1 cells
- Activated TH1 cells produce IFN- $\gamma$
- Important for response to intracellular infections
- Children born with deficient receptors have a weak Th1 response and low levels IFN- $\gamma$
- **Increased susceptibility:**
  - Disseminated Salmonella
  - Disseminated nontuberculous mycobacterial (NTM)
  - Disseminated Bacillus Calmette-Guerin (BCG) after vaccine
- **Treatment:** IFN- $\gamma$

# Penicillins

Jason Ryan, MD, MPH

## Penicillins




## Penicillin

Mechanism of Action

- Bacteria constantly breaking down/remaking cell wall
- Transpeptidases**
  - Cross link peptidoglycan in cell walls
  - Bind to **alanine** residues


## Penicillin

Mechanism of Action



## Penicillin


Mechanism of Action



## Penicillin

Mechanism of Action

- Penicillin binds to transpeptidases
  - "Penicillin binding proteins"
  - Mimics alanine ("D-alanyl-D-alanine") residues
  - Inactivates enzymes**
- Wall breakdown > wall creation → **Autolysis**
  - Enzymes that hydrolyze cell wall continue to work
  - Cell death (bactericidal)
- All  $\beta$ -lactam antibiotics: similar mechanism



## Penicillin

Mechanism of Action

Mestrother/Wikipedia

## Penicillin

Mechanism of Action

Alanine

D-Ala-D-Ala Terminus

Penicillin

Omarg10/Wikipedia

## Natural Penicillins

- Penicillin G (IM and IV)
- Penicillin VK (oral)
- **Probenecid**
  - Gout drug
  - Inhibits renal secretion PCN
  - Boosts PCN levels → co-administered in special circumstances

Penicillin G

Penicillin VK

## Natural Penicillins

Resistance

- **Modified penicillin binding proteins**
  - May result from genetic mutations
  - Example: *S. pneumonia* often produces altered PBPs
- **Reduced bacterial cell penetration**
  - Gram negative bacteria: poor penetration
  - Porins: gram negative proteins that transport nutrients/waste
  - Bacteria may decrease number of porins
- Beta lactamase enzyme

## Beta Lactamase

Penicillinase

- Bacterial enzymes
- Degrade beta lactam compounds
  - Penicillin G and VK
  - Some other penicillins
  - Some cephalosporins
- Many **gram negative bacteria**
- **Staphylococcus aureus**

## Beta Lactamase

- Most **gram negative rods** have beta-lactamase gene
  - "Chromosomal beta lactamase"
  - Some express low levels
- Genes can also be transferred via **plasmids**
  - Extra-chromosomal genetic material
  - Self-reproducing
  - Transferrable

## Beta Lactamase

- Gram negative bacteria
  - Beta lactamase found in **periplasm**
- Gram positive bacteria (*S. Aureus*)
  - No periplasm - Beta lactamase **secreted**
  - Generally produce more enzyme than GN




Image courtesy of Jeff Dahl/Wikipedia

## $\beta$ -Lactamase Inhibitors

Clavulanic Acid, Sulbactam, Tazobactam

- Inhibit bacterial  $\beta$ -lactamase
- Added to some penicillins to expand coverage
  - Aminopenicillins
  - Antistaphylococcal penicillins
- Little/no effect used alone

## Penicillin G and VK

Clinical Uses

- Narrow spectrum – few specific modern uses
- Gram positives
  - *Strep pyogenes* (strep throat)
  - *Actinomyces*
- *Treponema Pallidum* (syphilis)
- Rare uses (only in susceptible isolates)
  - *Neisseria meningitidis*
  - *Strep. pneumonia*

## Penicillin Adverse Effects

Hypersensitivity (allergic) reactions

- Commonly leads to hypersensitivity (allergic reaction)
- 1<sup>st</sup> exposure: Sensitization
- 2<sup>nd</sup> exposure: Hypersensitivity reaction
- Symptoms resolve on stopping drug

## Penicillin Adverse Effects

Hypersensitivity (allergic) reactions

- Acute (“immediate”)
  - **Type I, IgE-mediated**
  - Usually **within 1 hour** of taking drug
  - Histamine release
  - Itching, urticaria
  - Bronchospasm
  - Anaphylaxis



James Helman, MD

## Penicillin Adverse Effects

Maculopapular Rash

- “Non-immediate” reaction
- Most common with **aminopenicillins**
- Maculopapules
- Itchy or may be non-pruritic
- Absence of fever, wheezing, joint pain
- **Days or weeks** after starting drug
- **Type-IV (T-cell-mediated) mechanism**

Romano A et al. **Diagnosis of nonimmediate reactions to B-lactam antibiotics.** Allergy 2004

## Penicillin Adverse Effects

### Maculopapular Rash

- More common with viral infection
  - **EBV pharyngitis**
- Amoxicillin given for pharyngitis → maculopapular rash
- Mechanism not clear



Wikipedia/Public Domain

## Penicillin Adverse Effects

### Skin Reactions

- Stevens-Johnson Syndrome
  - Fever, necrosis
  - Sloughing of skin
  - Dermal-epidermal junction
  - Vesicles, blisters
- Toxic epidermal necrolysis
  - Severe form SJS (>30% skin)
- Mortality: SJS 1-5%; TEN 25-35%



Dr. Thomas Habif/Wikipedia

## Penicillin Adverse Effects

### Skin Reactions

- Immune mediated
  - CD8 T-cells play important roll
  - Re-challenge with drug can cause recurrence
- Antibiotic associations:
  - Sulfonamides (TMP-SMX)
  - **Aminopenicillins**
  - Cephalosporins

## Penicillin Adverse Effects

### Interstitial Nephritis

- Drug acts as hapten → immune response in kidneys
- Hypersensitivity (allergic) reaction
  - Complex mechanism
  - Considered a **Type IV hypersensitivity** reaction
  - T cells, Mast cells

Spanou Z et al. **Involvement of Drug-Specific T Cells in Acute Drug-Induced Interstitial Nephritis.** JASN Oct 2006

## Penicillin Adverse Effects

### Interstitial Nephritis

- Classic presentation
  - Fever
  - Oliguria
  - **Increased BUN/Cr**
  - **Eosinophils** in urine
  - White cells and **WBC casts** ("sterile pyuria")

## Penicillin Adverse Effects

### Hemolytic Anemia

- High doses can lead to extrinsic hemolytic anemia
- PCN binds to surface RBCs (hapten)
- Elicits immune response
  - **Antibodies against PCN bound to RBCs**
- Direct Coombs test: positive
- Type II hypersensitivity


## Penicillin Adverse Effects

Hypersensitivity (allergic) reactions

- Serum Sickness
  - Immune complex disorder (IgG)
  - **Days/weeks after exposure**
  - Complement activation
  - Type III hypersensitivity reaction
- Urticaria, fever, arthritis, lymphadenopathy

Tatum A et al. Severe serum sickness-like reaction to oral penicillin drugs: three case reports. Ann Allergy Asthma Immunol 2001

## Penicillin Immunology



## Penicillin Adverse Effects

C. Difficile Infection

- **Diarrhea** following antibiotic therapy
- Antibiotic depletes normal intestinal flora
- C. Difficile growth → **pseudomembranous colitis**
- May occur with any antibiotic
- Frequent associations
  - Clindamycin
  - Fluoroquinolones
  - Cephalosporins
  - **Penicillins**


## Jarisch-Herxheimer Reaction

- Occurs with **PCN therapy for spirochete infections**
- Classically occurs in **syphilis**
- Febrile syndrome
- Fever, chills, flushing, hyperventilation
- Usually ~2hrs after starting therapy
- Due to bacterial cell death → immune response

## Antistaphylococcal Penicillins

Oxacillin, nafcillin, dicloxacillin

- Side chain protects  $\beta$ -lactam from **staph penicillinase**
- Prototype: Methicillin
  - No longer used
  - High frequency of adverse effects (interstitial nephritis)
- Covers Staph Aureus (non-MRSA) and most strep



## Antistaphylococcal Penicillins



Oxacillin, nafcillin, dicloxacillin

- Common uses
  - Community acquired cellulitis
  - Impetigo
- Staph endocarditis based on culture data
- Side effects similar to penicillin



## Aminopenicillins

Amoxicillin/Ampicillin



## Aminopenicillins

Amoxicillin/Ampicillin

- Amoxicillin (oral)
- Ampicillin (IV)
  - Poor bioavailability when given orally
- **Penetrate porin channel of gram-negative bacteria**
- **Sensitive to beta lactamase enzymes**
- Covers penicillin bacteria plus some gram negatives

## Aminopenicillins

Amoxicillin/Ampicillin

### Bacteria

- **H. Influenza**
- E. Coli
- Proteus
- Salmonella
- Shigella
- **Listeria (gram +)**

### Main Clinical Uses

- Otitis Media
- Bacterial sinusitis
- Meningitis
  - Newborns, elderly
  - Listeria coverage

## Aminopenicillins

Maculopapular Rash

- Most common with aminopenicillins
- More common in viral infection
- Classic case
  - EBV infection with sore throat
  - Amoxicillin given for presumed bacterial pharyngitis
  - Maculopapular rash

## Aminopenicillins

Skin Reactions

- Stevens-Johnson Syndrome
- Toxic epidermal necrolysis
- Antibiotic associations:
  - Sulfonamides (TMP-SMX)
  - **Aminopenicillins**
  - Cephalosporins



Dr. Thomas Habib/Wikipedia

## $\beta$ -Lactamase Inhibitors


Clavulanic Acid, Sulbactam, Tazobactam

- Commonly used with aminopenicillins
  - Amoxicillin/Clavulanic acid (Augmentin)
  - Ampicillin/Sulbactam (Unasyn)
  - Increases activity against S. Aureus, H. flu
  - Also increases activity against anaerobes (B. fragilis)
- Common uses:
  - Otitis media/sinusitis (Broad-spectrum)
  - Bite wounds (Polymicrobial with anaerobes)


## Antipseudomonal Penicillins

Ticarcillin, Piperacillin

- **Greater porin channel penetration**
- Effective against *Pseudomonas aeruginosa*
- More gram (-) coverage vs. aminopenicillins



Ticarcillin  
(Carboxypenicillin)



Piperacillin  
(Piperazine penicillin)

## Antipseudomonal Penicillins


Ticarcillin, Piperacillin

- Susceptible to  $\beta$ -lactamases
- Given with  $\beta$ -lactamase inhibitor
  - Ticarcillin-clavulanate (Timentin)
  - Piperacillin-tazobactam (Zosyn)
- Broad-spectrum antibiotics
  - Most gram-positive (not MRSA)
  - More gram-negative (*pseudomonas*)
  - Most anaerobic bacteria
- Hospitalized patients with sepsis/PNA

# $\beta$ -Lactam Antibiotics

Jason Ryan, MD, MPH

## Penicillin Structure



## Beta Lactam Antibiotics


Carbapenems, Aztreonam, Cephalosporins

- Similar mechanism to penicillin
  - Bind transpeptidases (penicillin-binding proteins/PBPs)
  - Prevent peptidoglycan crosslinking
  - Autolysis
  - Usually **bactericidal**
- Potentially susceptible to beta lactamase

## Carbapenems

Imipenem, meropenem, ertapenem, doripenem

- $\beta$ -lactams (not penicillins)
- **Resistant to cleavage by most  $\beta$ -lactamase**



## ESBL

Extended Spectrum Beta Lactamase

- Plasmid-mediated bacterial enzymes
- Confer resistance to most beta-lactam antibiotics
  - Penicillins, cephalosporins, aztreonam
- Found only in gram-negative bacteria
  - Pseudomonas
  - Klebsiella
  - E. coli
  - Enterobacter
  - Salmonella
  - Serratia
  - Shigella

## Carbapenems

Imipenem, meropenem, ertapenem, doripenem

- Drug of choice for ESBL bacteria
- Broad spectrum:
  - Gram (+)
  - Gram (-) including pseudomonas, enterobacter
  - Anaerobes including B. fragilis
- Used in hospitalized patients

## Imipenem

- First commercially available carbapenem
- Metabolized in kidneys
  - Loss of antibacterial effect
  - Nephrotoxic metabolites
- Proximal tubule enzyme: **dehydropeptidase I**
- Given with **cilastatin** (enzyme inhibitor)

## Carbapenems

Imipenem, meropenem, ertapenem, doripenem

- Imipenem and meropenem
  - Older carbapenems
  - No important differences in efficacy
- Doripenem and ertapenem
  - Newer carbapenems
  - Doripenem: Similar to imipenem and meropenem
  - Ertapenem: Some resistance in ESBL bacteria
- Ertapenem
  - Once daily dosing
  - Weak activity against pseudomonas


## Carbapenems

Imipenem, meropenem, ertapenem, doripenem

- Common side effects
  - Nausea, vomiting, diarrhea
  - Skin rash
- Neurotoxicity
  - Seizures
  - Inhibition of GABA receptors
  - Especially at high doses or with renal failure
  - Lower risk with **meropenem**

## Aztreonam

- Monobactam:  $\beta$ -lactam ring not fused to another ring





## Aztreonam

- Binds penicillin-binding protein 3 (PBP-3)
  - Found in **gram negative bacteria**
  - Prevents cross-linking of peptidoglycan
  - Bactericidal
- Limited susceptibility to  $\beta$ -lactamase
  - Some resistance in ESBL bacteria
- Only active against **gram (-) bacteria**
  - Does not bind PBP of gram (+) bacteria
  - No activity against gram (+) or anaerobes
  - Active against **pseudomonas**

## Aztreonam

- Intravenous administration (hospitalized patients)
- Synergistic with **aminoglycosides**
- No cross reactivity in **penicillin allergic patients**
- Key niche: penicillin allergy

## Cephalosporins



Images courtesy of Wikipedia/Public Domain

## Cephalosporins

- Divided into 1<sup>st</sup> through 4<sup>th</sup> generation
- 1<sup>st</sup> generation: Mostly gram positive coverage
- Successive generations: increased gram (-) coverage

## 1<sup>st</sup> Generation Cephalosporins

- Cefazolin, cephalexin
- Developed to treat S. Aureus resistance to penicillin
- Covers many gram (+) including S. Aureus (not MRSA)
  - **Stable against S. Aureus beta lactamase**
  - Does not cover enterococcus or listeria
  - Susceptible to gram negative beta lactamases
- Main uses:
  - Surgical wound (skin) infections
  - Cefazolin given pre-op for prevention

## 2<sup>nd</sup> Generation Cephalosporins

Cefuroxime, Cefoxitin, Cefotetan

- Developed to treat amoxicillin-resistant infections
- Increased affinity for gram (-) PBPs
- **More resistant to beta lactamase**
- Increased gram (-)
  - H. influenza, Enterobacter, Proteus
  - E. coli, Klebsiella, Serratia, N. gonorrhoea
- Increased anaerobic coverage (B. fragilis)

## 2<sup>nd</sup> Generation Cephalosporins

Cefuroxime, Cefoxitin, Cefotetan

- Cefuroxime (oral):
  - Otitis media (S. pneumonia, H. flu)
  - UTI in children (E. coli; no fluoroquinolones)
- Cefoxitin/cefotetan (IV):
  - PID (covers Neisseria; also give doxycycline for Chlamydia)
  - Pre-op in children with appendicitis
    - E. coli
    - Covers gram negatives and some anaerobes
    - Usually given with metronidazole

## 3<sup>rd</sup> Generation Cephalosporins

Ceftriaxone, Cefotaxime, Ceftazidime

- Broad gram (-) coverage
  - **More resistance to beta lactamase enzymes**
  - More gram (-) PBP affinity
- Ceftriaxone, Cefotaxime: Poor coverage pseudomonas
- Ceftazidime: Covers pseudomonas
  - Used in hospitalized patients with gram negative infections
  - Sepsis/pneumonia
- Most achieve **good CSF penetration** (meningitis)

## Ceftriaxone

- Commonly used for N.gonorrhoea
- Commonly used in meningitis
  - Active against S. pneumoniae, N. meningitidis
  - Good CSF penetration


## 4<sup>th</sup> Generation Cephalosporins

Cefepime

- Broad spectrum (>3<sup>rd</sup> generation drugs)
  - MSSA
  - Many gram (+)'s
  - Many gram (-)'s including **pseudomonas**
- Resistant to some ESBL
- Hospitalized patients with gram (-) infections

## $\beta$ -lactamase Sensitivity

Based on side chain



- \* Anti-staphylococcal penicillins resist staph penicillinase  
\*\* Gram negatives only

## 5<sup>th</sup> Generation Cephalosporins

Ceftaroline

- Active against **MRSA**
- FDA approval 2010
- Prodrug converted to active metabolite
- Binds **PPB2a**
  - MRSA-specific PBP
  - Low affinity for most other beta-lactams
- Covers MRSA and VRSA
- Some gram negatives (not pseudomonas)
- Studied in skin infections and pneumonia

## Cephalosporins

Resistance Mechanism

- Modified penicillin-binding proteins (PBPs)
- Altered cell permeability
- Beta lactamase

## Cephalosporins

Adverse Reactions

- **Hypersensitivity Reactions** (similar to PCN)
  - Anaphylaxis
  - Maculopapular rash
  - Serum sickness (fever, rash, arthritis)
  - Hemolytic anemia (drug as hapten)
  - Interstitial nephritis
  - Stevens-Johnson Syndrome/Toxic epidermal necrolysis
- Some cross-reactivity with penicillins
  - Traditionally cited as 10%
  - Actual risk may be lower


## Cephalosporins

### Adverse Reactions

- **Vitamin K deficiency**
  - Vitamin K1 from diet (green, leafy vegetables)
  - Vitamin K2 from GI bacteria
  - Antibiotics reduce bacterial vitamin K production
  - Result: Increased INR and potential bleeding
  - Commonly a problem for **patients on warfarin**
  - May be caused by any antibiotic

## Cephalosporins

### Adverse Reactions



- **Hypoprothrombinemia**
  - Associated with N-methylthiotetrazole (NMTT) side chains
  - Cefotetan, cefazolin
  - Inhibits epoxide reductase (similar to warfarin)
  - ↓ hepatic synthesis of clotting factors
  - May prolong the PT/INR
  - Reversible with vitamin K
  - Most reports among malnourished patients

Shearer et al. Mechanism of cephalosporin-induced hypoprothrombinemia: relation to cephalosporin side chain, vitamin K metabolism, and vitamin K status. J Clin Pharmacol. 1988

## Cephalosporins

### Adverse Reactions


- Nephrotoxicity of aminoglycosides
  - Reports of increased risk with combination therapy

## Cephalosporins

### Adverse Reactions

- **Disulfiram reaction**
- Alcohol consumption with cephalosporins
- Warmth, flushing, sweating
- Inhibition of acetaldehyde dehydrogenase
- Accumulation of acetaldehyde
- Occurs with certain side chain structures
- Cefoperazone, cefamandole, and cefotetan

## Ethanol Metabolism




# Sulfonamides

Jason Ryan, MD, MPH


## Sulfonamide Antibiotics

- Sulfonamide group =  $\text{SO}_2\text{-N}$
- "Sulfa" drug = Contains sulfonamide group




## Bacterial Folate Synthesis


- Folate required for thymidine/DNA synthesis
- Mammalian cells: use exogenous folate (diet)
- Bacterial cells: no exogenous folate (must synthesize)




## Bacterial Folate Synthesis



## Bacterial Folate Synthesis




## Bacterial Folate Synthesis



## Sulfonamides

Sulfamethoxazole (SMX), sulfisoxazole, sulfadiazine

- Mimics of PABA
- Competitively inhibit dihydropteroate synthase




## Sulfonamides

Resistance

- Increased PABA
- Altered dihydropteroate synthase
- Decreased uptake

## Dapsone

- Not a sulfonamide
- Competes with PABA for dihydropteroate synthase
- Two main uses:
  - Mycobacterium leprae (leprosy)
  - Pneumocystis jirovecii



## Sulfonamides


Sulfamethoxazole (SMX), sulfisoxazole, sulfadiazine

- Usually given with trimethoprim
  - TMP-SMX (Bactrim)
  - Sequential block of THF synthesis
- Sulfadiazine
  - Silver-sulfadiazine (cream) for burns
- Sulfadiazine and pyrimethamine
  - Also sequential block of THF synthesis
  - Used in toxoplasmosis (HIV)

## Sulfonamides

Toxicity


- Hypersensitivity reactions ~3% of patients
- Reactions linked to:
  - Arylamine (NH<sub>2</sub>) at N4 position
  - Nitrogen ring attached to N1 nitrogen
- Only sulfonamide antibiotics contain both features



## Sulfonamides

Toxicity

- Other sulfa drugs



## Sulfonamides

Hypersensitivity (allergic) reactions

- Similar to penicillin allergic reactions
- Anaphylaxis
- Maculopapular rash
- Serum sickness (fever, rash, arthritis)
- Interstitial nephritis
- **Stevens-Johnson Syndrome**
- **Toxic epidermal necrolysis**

## Sulfonamides

Toxicity

- Photosensitivity
- Drug interaction with UV light
- Caused by many drugs
- Common drugs
  - Tetracycline
  - **Sulfonamides**
  - Amiodarone



Wikipedia/Public Domain

## Sulfonamides

Toxicity

- Hemolysis in **G6PD deficient** patients
  - RBC susceptible to oxidative stress
  - Sulfonamides are oxidants - classic trigger for hemolysis
  - Other triggers: **Dapsone**



Database Center for Life Science (DBCLS)

## Sulfonamides

Toxicity

- Binds to **albumin**
- Displaces other bound substances
  - Bilirubin
  - Warfarin

## Sulfonamides

Toxicity

- **Kernicterus** in infants
  - Sulfonamides → increased free bilirubin levels
  - Unconjugated bilirubin: neurotoxic
  - Basal ganglia, brainstem nuclei
- Permanent neurologic impairment
  - Movement disorder (chorea, tremor)
  - Hearing loss
  - Limited gaze

## Sulfonamides


Toxicity

- Raise **warfarin** levels
  - Displaces warfarin from albumin
- INR level may rise in patients on warfarin therapy

## Trimethoprim/Pyrimethamine

### Mechanism of Action

- Inhibit **dihydrofolate reductase**
- Similar structure to dihydrofolate



## Trimethoprim/Pyrimethamine

### Toxicity

- Preferentially inhibits bacterial DHF reductase
- Some inhibition of human enzyme can occur
- Inhibits DNA synthesis of rapidly dividing cells


## Trimethoprim/Pyrimethamine

### Toxicity

- Bone marrow suppression
  - Pancytopenia: megaloblastic anemia, leukopenia, ↓platelets
- Can alleviate with **leucovorin (folinic acid)**
  - Converted to THF
  - Does not require dihydrofolate reductase
  - "Leucovorin rescue"

## Dihydrofolate Reductase

### Human DNA Synthesis



## TMP-SMX

### Bactrim

- Combination is **bactericidal**
- Covers many gram (+) and gram (-)
  - Does not cover pseudomonas
  - Does not cover B. fragilis (and most anaerobes)
- Covers some fungi and parasites
- Common uses:
  - Urinary tract infections (covers E. Coli well)
  - Pneumocystis pneumonia in HIV (treatment/prophylaxis)

## TMP-SMX

### Pregnancy

- Risk of **kernicterus**
- Disrupts **follic acid metabolism**

## PCP

*Pneumocystis jirovecii*

- Opportunistic fungal infection
- Occurs in end stage HIV/AIDS
- Treatment of choice: TMP-SMX
- Hypersensitivity reactions: **6-25x higher in HIV**
- Alternative therapy often needed
  - Dapsone
  - Pentamidine
  - Atovaquone (malaria drug)




Image courtesy of Yale Rosen/Wikipedia

# Protein Synthesis Inhibitors

Jason Ryan, MD, MPH


## Protein Synthesis Inhibitors

- Aminoglycosides
- Macrolides
- Tetracyclines
- Chloramphenicol
- Clindamycin
- Linezolid
- Streptogramins

## Bacterial Protein Synthesis

- DNA → Transcription → RNA
  - RNA polymerase
  - Target of Rifampin (tuberculosis)
- RNA → Translation → Protein

## Bacterial Protein Synthesis



## Aminoglycosides

Gentamicin, Neomycin, Amikacin, Tobramycin, Streptomycin

- Block initiation of protein synthesis
  - Primarily bind 30S
  - Misreading of genetic code
  - Bacteria cannot divide, produce cellular proteins
  - Cell death (bactericidal)
- Require O<sub>2</sub> for transport into cells
  - Not effective against anaerobes
- Do not effect eukaryotic cells (different ribosomes)
- Not transferred into eukaryotic cells
  - Not effective intracellular organisms (Rickettsia/Chlamydia)

## Aminoglycosides

Gentamicin, Neomycin, Amikacin, Tobramycin, Streptomycin

- Used alone (rare) to treat serious gram (-) infections
- Streptomycin can be used for tuberculosis
  - Older agent
  - Used in combination with other drugs
- Neomycin given prior to bowel surgery
  - Poorly absorbed (stays in gut)
  - Often given with Erythromycin
  - Decrease colonic bacteria

## Aminoglycosides

Gentamicin, Neomycin, Amikacin, Tobramycin, Streptomycin

- Often added to  $\beta$ -lactams
  - Synergistic effects
  - Combination more effective than sum of effects
- Vancomycin/gentamicin for endocarditis
- Ampicillin/gentamicin for newborn meningitis
- Pip/Tazo + tobramycin for CF patients (pseudomonas)

## Aminoglycosides

Resistance

- Most common mechanism resistance:
  - "Aminoglycoside modifying enzymes"
  - Bacteria acquire enzymes that modify drug structure
  - Modified structure binds poorly to ribosomes
  - Phosphorylation (mediated by aminoglycoside kinases)
  - Adenylation/acetylation (mediated by transferases)

## Aminoglycosides

Adverse Effects

- Ototoxicity
  - Toxic to 8<sup>th</sup> cranial nerve
  - Hearing loss, balance problems (falls)
  - Mechanism not clear
- Nephrotoxicity
  - Acute tubular necrosis
  - 5-10% of drug taken up by proximal tubular cells
  - Serum Cr will rise

## Aminoglycosides

Adverse Effects

- Neuromuscular blockade
  - Rare side effect
  - Can block/limit release of ACh at neuromuscular junctions
  - Usually occurs when levels are high or pre-existing neuromuscular disease
- Pregnancy class D
  - Reports of renal and ototoxicity in fetus

## Aminoglycosides

Monitoring

- Plasma levels sometimes monitored
  - Trough level: Just before next dose
  - Peak level: Short time after dose
- High trough = risk of toxicity
- Low peak = less effective therapy

## Macrolides

Azithromycin, Clarithromycin, Erythromycin

- Block translocation
- 50S ribosomal subunit
- Covers many gram (+) cocci, especially strep
- Some gram (-) coverage
- Concentrated inside macrophages, other cells
- Effective against intracellular pathogens
  - Chlamydia (obligate), Legionella (facultative)

## Macrolides

Azithromycin, Clarithromycin, Erythromycin

- Community acquired pneumonia
  - Azithromycin covers Strep, H. flu, Atypicals
  - Good for penicillin allergic patients
- Chlamydia infection
  - Azithromycin (safe in pregnancy)
  - Often co-administered with Ceftriaxone (gonorrhea)

## Macrolides

Azithromycin, Clarithromycin, Erythromycin

- Erythromycin
  - Binds to motilin receptors in GI tract
  - Stimulates smooth muscle contraction
  - Can be used in GI motility disorders
- Clarithromycin
  - Part of triple therapy for H. pylori

## Macrolides

Resistance

- Resistance mechanism
  - 23S rRNA = component of 50S ribosome
  - Location of macrolide binding
  - Methylation of this site → resistance

## Macrolides

Adverse Effects

- Nausea, diarrhea, abdominal pain (motility)
  - Erythromycin worst offender
- Prolonged QT on EKG
  - Erythromycin also worst offender
- Acute cholestatic hepatitis
  - ↑AST/ALT/Alk Phos/Bilirubin
  - Case reports in patients on Azithromycin
  - Contraindicated with history of cholestatic jaundice or hepatic dysfunction

## Macrolides

Adverse Effects

- Rash
  - Maculopapular allergic reaction
- P450 Enzyme Inhibitors
  - Will raise serum levels of P450 metabolized drugs
  - Theophylline, Warfarin

## Tetracyclines

Tetracycline, doxycycline, demeclocycline, minocycline

- Transported into bacterial cells
- Binds 30S ribosome
- Prevents attachment of tRNA
- Demeclocycline
  - Not used as an antibiotic
  - ADH antagonist
  - Given in SIADH
  - Differentiates Central from Nephrogenic DI

## Doxycycline

- Most commonly used member tetracycline family
- Accumulates intracellularly
- Covers many unusual/atypical bacteria
  - Most zoonoses
  - Chlamydia
- Used to treat acne vulgaris (also minocycline)
  - Covers propionibacterium acnes within follicles

## Tetracyclines

Tetracycline, doxycycline, demeclocycline, minocycline

- Absorption impaired by minerals and antacids
  - Calcium, magnesium (antacids)
  - Iron
  - Dairy including milk
- These substances are cations that chelate the drug
- Cannot be taken with antacids or milk

## Tetracyclines

Resistance

- Decreasing influx or increasing efflux from cells
- Plasmid-encoded transport pumps
- Different from many other antibiotics:
  - No alteration of drug by bacteria

## Tetracyclines

Adverse Effects

- GI distress (common)
  - Epigastric pain, nausea, vomiting and anorexia
- Photosensitivity
  - Red rash or blisters in sun exposed areas

## Tetracyclines

Adverse Effects

- Discoloration of teeth
  - Brown-yellow discoloration of teeth
  - Children under the age of eight (does not occur in adults)
- Inhibition of bone growth in children
  - Deposit in bones
  - Chelate with calcium
- Contraindicated in pregnancy
  - Cross placenta
  - Can accumulate in fetal bone and teeth

## Chloramphenicol

- 50S ribosome
- Inhibits peptidyl transferase
- Rarely used in developed world:
  - Toxicity
  - Increasing resistance
- Used in developing world due to low cost

## Chloramphenicol

- Broad coverage of gram (+), gram (-), atypicals
- Can be used in pregnancy instead of doxycycline
  - Rickettsia (RMSF), Ehrlichia
  - Only in 1<sup>st</sup>/2<sup>nd</sup> trimester
  - 3<sup>rd</sup> trimester risk of gray baby syndrome
- Can be used for meningitis (developing world)
  - Covers Neisseria
  - Less effective than alternative drugs

## Chloramphenicol

### Adverse effects

- Anemia
  - Bone marrow suppression
- Aplastic anemia
  - Idiosyncratic
  - Irreversible → often fatal
- Gray baby syndrome
  - Babies lack liver UDP-glucuronyl transferase
  - Required for metabolism/excretion of drug
  - Skin turns ashen, gray
  - Hypotension
  - Often fatal

## Clindamycin

- 50S ribosome
  - 23S rRNA component
  - Same as macrolides
- Covers some gram (+)
  - Staph, viridans strep, Strep pyogenes, and S. pneumoniae
- Covers many anaerobes
  - Clostridium perfringens
  - Mouth anaerobes: Fusobacterium, Prevotella, Peptostreptococcus

## Clindamycin

- Main use is to cover anaerobes “above the diaphragm”
  - Aspiration pneumonia
  - Lung abscesses
  - Oral infections (mouth anaerobes)
- Lots of resistance to clindamycin in B. fragilis
  - Anaerobic infections “below the diaphragm”
  - Metronidazole

## Clindamycin

### Adverse Events

- Classic cause of C. difficile infection
  - Up to 10% of patients
  - Pseudomembranous colitis
  - C. difficile overgrowth
  - Massive, watery diarrhea
- Antibiotic-associated diarrhea
  - Milder than C. diff infection
  - Changes in GI flora
  - Less absorption of solutes → osmotic diarrhea
  - Stops when drug discontinued

## Linezolid

- Binds to 50S Ribosome
- Blocks initiation
- Main use: Vancomycin-resistant enterococcus (VRE)
  - Epidemics in hospitals
  - Usually occurs in patients with prior antibiotic treatment

## Linezolid

- Weak monoamine oxidase (MAO) inhibitor
- Can cause serotonin syndrome
- High risk when given with SSRIs
- Fever, confusion, agitation, hyperreflexia

## Streptogramins

quinupristin/dalfopristin

- Block protein synthesis 50S ribosome
- Used together for sequential protein synthesis block
  - Synercid (quinupristin/dalfopristin)
- Used for vancomycin resistant bacteria
  - VRSA
  - VRE

## Bacteriostatic vs. Bactericidal

Antibiotic Class	Action
Aminoglycosides	Bactericidal
Macrolides	Bacteriostatic
Tetracyclines	Bacteriostatic
Chloramphenicol	Bacteriostatic
Clindamycin	Bacteriostatic
Linezolid	Bacteriostatic (mostly)
Quinupristin/dalfopristin	Variable

Most protein synthesis inhibitors are bacteriostatic  
Only aminoglycosides are bactericidal  
Misread proteins travel to membrane and increase permeability

Source: Microbiol Rev. Sep 1987; 51(3): 341-350.

## Other Antibiotics

Jason Ryan, MD, MPH


## Other Antibiotics

- Quinolones
- Vancomycin
- Metronidazole
- Nitrofurantoin

## Quinolones

Ciprofloxacin, Levofloxacin, Moxifloxacin, Norfloxacin

- Inhibit enzymes for bacterial **DNA synthesis**
- **DNA gyrase**
- **Topoisomerase IV**



## Quinolones

Ciprofloxacin, Levofloxacin, Moxifloxacin, Norfloxacin

- Bacterial topoisomerase enzymes
- DNA Gyrase
  - Introduces double-stranded break
  - Repairs break
- Topoisomerase IV
  - Separates daughter chromosomes
  - "Decatenation"
- Inhibition → DNA damage → **cell death**

## Quinolones

Resistance Mechanisms

- Alterations of **DNA gyrase** and **topoisomerase IV**
- Alteration in cell permeability
- Efflux of drug

## Quinolones


Ciprofloxacin, Levofloxacin, Moxifloxacin, Norfloxacin

- Many gram (+), gram (-), atypicals
- Common clinical uses (adults only)
  - UTIs (E. Coli, other enteric gram negatives)
  - Pneumonia (S. pneumo, H. flu, atypicals)
  - Abdominal infections (enteric gram negatives)

## Quinolones

### Early Drugs

- Nalidixic acid (not a fluoroquinolone), Norfloxacin
- Mostly gram negative coverage
- Limited/no gram positive coverage




Nalidixic Acid

## Quinolones

### Ciprofloxacin


- Some gram positive coverage
  - Rarely used alone for gram positive coverage (resistance)
- Very good **gram negative coverage**
- Most reliable **pseudomonas** coverage
- Used in UTIs, GI infections
- Cipro ear drops for otitis externa



## Quinolones

### Levofloxacin

- More gram positive/atypical coverage than Cipro
  - Better strep pneumo coverage than Cipro
  - Covers most methicillin-susceptible Staph aureus
- Less effective against pseudomonas than Cipro
- Commonly used in pneumonia (strep, atypicals)



## Quinolones

### Gatifloxacin, Sparfloxacin, Moxifloxacin

- Better gram (+)/atypical coverage than Levofloxacin
- Less effective for pseudomonas than Levofloxacin
- Also used in pneumonia

## Quinolones


### Adverse Reactions

- **Gastrointestinal** upset
  - Anorexia, nausea, vomiting, and abdominal discomfort
  - Up to 17% of patients
- **Neurologic** side effects
  - Headache, dizziness
  - 2 to 6% of patients

## Quinolones

### Adverse Reactions

- QT prolongation on EKG
- Caused by blockade of K<sup>+</sup> channels
- Can lead to torsade de pointes



## Quinolones

### Adverse Reactions

- **Tendon rupture/tendonitis**
  - Most commonly Achilles
  - More common older patients (>60), people on steroids
- Cannot use in pregnancy/children
  - Toxic to developing cartilage in animal studies




## Antacids

- Disrupt absorption of many drugs
- Aluminum and magnesium hydroxide
- Sucralfate (contains aluminum)
- Key drugs
  - Tetracycline
  - **Fluoroquinolones**
  - Isoniazid
  - Iron supplements

## Vancomycin

- Inhibits peptidoglycan formation (cell wall)
- Binds **D-alanyl-D-alanine peptides**
- Prevents crosslinking
- Cell wall breakdown > formation → **cell death**
- Same effect as beta lactams via different mechanism
  - Beta lactams: inhibit transpeptidases
  - Vancomycin: block transpeptidase binding


## Vancomycin




Mcstrother/Wikipedia

## Vancomycin

- Resistance: terminal amino acids change
  - D-alanyl-D-alanine changed to **D-alanyl-D-lactate**
  - VRSA emerges




Alanine



Lactate

## Vancomycin

- Only effective in gram (+)
- Too large to pass outer membrane gram negatives



## Vancomycin

- Two common uses:
  - #1: Methicillin resistant Staph Aureus (MRSA)
  - #2: Oral therapy for *C. difficile* pseudomembranous colitis
- Often given empirically when MRSA is a concern
  - Endocarditis
  - Severe pneumonia/sepsis

## Vancomycin

### Adverse Effects

- Generally well tolerated
- Nephrotoxicity
  - Less common with modern preparations
  - Increased risk if **concomitant aminoglycoside therapy**
- Ototoxicity
  - Tinnitus, vertigo, and hearing loss reported (rare)


## Vancomycin

### Adverse Effects

- **Red man syndrome**
  - Flushing, erythema, itching
  - Usually affects upper body, neck, face more than lower body
  - Occurs 10-20 minutes after start of infusion
- Direct activation of mast cells → **histamine** release
  - "Pseudoallergic drug reaction"
- May develop with first administration
- Infusion related → slow infusion = no symptoms

## Metronidazole

- Prodrug: Must be **reduced** to activate
- Only **anaerobic bacteria** capable of reduction
- Reduced metronidazole → more drug uptake
- Activated form generates free radicals
- Interact with DNA
- DNA breakage/destabilization
- Cell death



## Metronidazole

### Uses

- Good coverage of anaerobes "below the diaphragm"
  - **Bacteroides fragilis**
  - **Clostridium difficile**
- Peritonitis, abdominal abscesses, diverticulitis
- Often given with quinolone for anaerobic/GI gram(-)
- Cipro/Flagyl often used for diverticulitis

## Metronidazole

### Uses

- **H. pylori** and **Gardnerella vaginalis**
  - Facultative anaerobic bacteria
  - Susceptible to metronidazole
- Triple therapy for H. Pylori
- Treatment of bacterial vaginitis

## Metronidazole

### Uses

- Anaerobic protozoa (lack mitochondria)
  - **Trichomonas vaginalis**
  - **Entamoeba histolytica**
  - **Giardia lamblia**
- Covered by metronidazole

## Metronidazole

### Adverse Reactions

- Unpleasant metallic taste
- **GI**: Abdominal discomfort, nausea
- **Neuro**: Neuropathy, headache

## Metronidazole

### Adverse Reactions

- **Disulfiram-like reaction**
- Alcohol consumption with metronidazole
- Warmth, flushing, sweating
- Unclear mechanism
- Metronidazole may not inhibit alcohol metabolism
- Patients should **avoid alcohol**



Visapää JP. Lack of disulfiram-like reaction with metronidazole and ethanol. *Ann Pharmacother.* 2002. Jun;36(6):971-4.

## Nitrofurantoin

- Rarely used antibiotic
- Exact mechanism incompletely understood
- Bactericidal drug
- Only use is **UTIs** (concentrates in urine)
- Two things to know about this drug:
  - Used for UTIs in **pregnancy** (avoid TMP-SMX, quinolones)
  - Can trigger **hemolysis in G6PD patients**

# Fungal Pneumonias

Jason Ryan, MD, MPH

## Fungi

- Fungi are eukaryotes
  - Have a nucleus
  - Intracellular organelles
  - VERY different from bacteria
- Single celled: yeast
- Multicellular
  - Filamentous molds
  - Mushrooms
- Dimorphic = yeast or filamentous

## Fungi

- Hyphae: long, branches of fungi
  - Often divided into multiple cells by septa (septate hyphae)
- Mycelium
  - Many hyphae together
- Spores (conidia)
  - Often formed on ends of hyphae (conidiophore)




Image courtesy of Y Tambe/Wikipedia



Image courtesy of Wikipedia/Public Domain

## Fungal Infections

- Pneumonia
  - Only in specific geographic areas
  - Histoplasmosis, Blastomycosis, Coccidioidomycosis
- Skin
  - Tinea versicolor, tinea pedis, sporothrix
- Opportunistic
  - Candida, Aspergillus, Cryptococcus, Mucormycosis, Pneumocystis

## Pneumonia Fungal Infections

- Histoplasmosis
- Blastomycosis
- Coccidioidomycosis
- Paracoccidioidomycosis

## Pneumonia Fungal Infections

### Key features

- Cause pneumonia but can disseminate
- Dimorphic except coccidioidomycosis
- Cold temps = mold; warm temps (body) = yeast
- Disease from inhaling fungus
  - Not person to person spread
- All can cause granulomatous inflammation
  - Potential TB mimics
- Treatment:
  - Mild disease: Fluconazole/itraconazole
  - Severe/systemic: Amphotericin B

## Granulomatous Infections

- Tuberculosis, Leprosy
- Fungal pneumonias (Histo, Blasto, Coccidio)
- Bartonella (cat scratch disease)
- Brucella
- Listeria in infants (Granulomatosis Infantiseptica)
- Schistosomiasis (worm)
- Syphilis (gummas)

## Pneumonia Fungal Infections

- All have two distinct clues to diagnosis
  - Geography
  - Pathology

## Geography

- Histoplasmosis
  - Ohio and Mississippi river valleys
- Blastomycosis
  - Ohio and Mississippi river valleys
  - Great Lakes
- Coccidioidomycosis
  - Southwest US
- Paracoccidioidomycosis
  - South/Central America

## US Map



## Geography

- Histoplasmosis/Blastomycosis
  - Ohio River/valley
  - Mississippi River/valley
  - "Midwest"
  - Great Lakes (Blastomycosis)
- Coccidioidomycosis
  - Arizona, New Mexico, California
- Paracoccidioidomycosis
  - South/Central America

## Histoplasmosis

*Histoplasma capsulatum*



- Soil contaminated with bird or bat droppings
- Key site for infection: Caves
- Spores inhaled → yeast at body temperature
- Ingested by macrophages
- Survive/multiply in macrophages
  - Can be spread to others

## Histoplasmosis

### Symptoms/Treatment

- Vast majority of people asymptomatic
- Most common symptoms are pulmonary
  - Slow onset over weeks
  - Mild pneumonia
  - Hilar lymphadenopathy
- Diagnosis: Antigen/antibody tests, biopsy
- Treatment:
  - Mild disease: Fluconazole/itraconazole
  - Severe/systemic: Amphotericin B

## Histoplasmosis



Images courtesy of Nivaldo Medeiros, MD, [www.hematologyatlas.com](http://www.hematologyatlas.com). Used with permission

## Histoplasmosis

### Disseminated disease

- Cell-mediated immunity clears infection
- Dissemination rare unless immunocompromised
  - HIV/AIDS
  - TNF- $\alpha$  inhibitors
- LOTS of symptoms: GI, CNS, anemia, lymph
- Some key features
  - Hepatosplenomegaly (abnormal LFTs)
  - Tongue, mouth ulcers
  - Pancytopenia (bone marrow involvement)
- Treatment is Amphotericin B

## Blastomycosis

### Blastomyces dermatitidis

- Inhaled conidia  $\rightarrow$  yeast in the body (dimorphic)
- Many patients asymptomatic
- When symptomatic: slow onset PNA most common

## Blastomycosis

- Extrapulmonary disease in ~20% of patients
- Skin is most common site
  - Verrucous (warts) lesions with irregular borders
- Bone (osteomyelitis) next most common
- Classic case:
  - Mississippi river exposure
  - Slow onset pneumonia
  - Skin lesions
  - Possibly bone pain

## Blastomycosis

- Forms granulomas
- Classic path finding is "broad based budding yeast"
- Diagnosis by visualizing yeast
  - Sputum
  - Tissue
- Treatment:
  - Mild disease: Fluconazole/itraconazole
  - Severe/systemic: Amphotericin B




Image courtesy of Medmyco/Wikipedia

## Coccidioidomycosis

- Mexico, Arizona, New Mexico, Texas, California
- Grows as mold beneath desert surface
- Dry conditions → Mold fractures into spores
- Infection by inhalation of a spores
- In lung, spore enlarges to spherule (not a yeast)
- Enlarging spherules produce endospores
- Mature spherules rupture, releasing endospores
- Each endospore can produce another spherule

## Spherule

- Classic path finding: Spherule filled with endospores




Image courtesy of Wikipedia/Public Domain

## Coccidioidomycosis


- Most infections asymptomatic
- Symptoms: Valley Fever
  - Fever, malaise, cough
  - Arthralgias
  - Erythema nodosum
- Diagnosis: sputum Cx
- Rarely disseminates
- Feared result is meningitis
- Treatment:
  - Mild disease: Fluconazole/itraconazole
  - Severe/systemic: Amphotericin B



Image courtesy of James Helman, MD

## Paracoccidioidomycosis

- Central/South America
- Pulmonary symptoms (cough)
- In yeast form, mother cells buds off children
  - "Pilot's wheel"
  - "Mickey Mouse Head"



CDC (Dr. Lucille K. George)/Public Domain



Image courtesy of Wikipedia/Public Domain

## Aspergillus

- Pneumonia in immunocompromised
  - HIV, Chemotherapy
- No specific geographic area
- Very sick patient
  - Fever, hemoptysis, pleuritic chest pain

# Fungal Skin Infections

Jason Ryan, MD, MPH

## Fungal Infections

- Pneumonia
  - Only in specific geographic areas
  - Histoplasmosis, Blastomycosis, Coccidioidomycosis
- Skin
  - Tinea versicolor, tinea pedis, sporothrix
- Opportunistic
  - Candida, Aspergillus, Cryptococcus, Mucormycosis, Pneumocystis

## Skin and Nail Infections

- Dermatophyte infections
  - Tinea pedis
  - Tinea cruris
  - Tinea corporis
  - Tinea capitis
  - Tinea unguium
- Tinea versicolor
- Sporothrix schenckii

## Terminology

- Tinea = fungal skin/nail infection
- Dermatophytes: fungi that require keratin for growth
- Most tinea infections caused by dermatophytes

## Dermatophytes

- Majority of infections from 3 dermatophytes:
  - Epidermophyton
  - Trichophyton
  - Microsporum
- All consume keratin
- Exist only as molds with hyphae
- Most treated with topical antifungals
  - Clotrimazole, Miconazole (azoles)
  - Terbinafine
- Note: Nystatin not effective against dermatophytes
  - Only effective against cutaneous candida (diaper rash)

## KOH Prep

- Potassium hydroxide (KOH)
- Used to identify fungal infections
- KOH dissolves epidermal keratinocytes
- Fungi visible in skin scrapings (hyphae)
- Used for:
  - Dermatophyte infections
  - Tinea versicolor
  - Candida

## Tinea unguium

Onychomycosis

- Mostly a cosmetic problem
- Oral treatment often used:
  - Terbinafine
  - Itraconazole



Image courtesy of James Hellman/Wikipedia

## Tinea pedis

Athlete's foot

- Fungal foot infection
- Itchy, red erosions between toes, on soles
- Untreated can lead to scaling
- Treatment: Topical antifungals
  - Clotrimazole, Miconazole
  - Terbinafine

## Tinea corporis

Ringworm

- Itchy, circular or oval, red, scaling patch or plaque
- Spreads centrifugally
- Red border with central clearing ("ring")
- Treatment: Topical antifungals
  - Clotrimazole, Miconazole
  - Terbinafine



Image courtesy of Dr. Lucille K. Georg/Wikipedia

## Tinea capitis

- Dermatophyte infection of scalp
- Usually occurs in children
- Common in African-Americans
- Red, scaling patch on the scalp
- Spreads centrifugally
- Oral treatment often used:
  - Griseofulvin
  - Terbinafine
  - Itraconazole



Image courtesy of myself/Wikipedia

## Tinea cruris

Jock itch

- More common in men
- Often occurs after physical activity with sweating
- Obesity increases risk
- Red patch on inner thigh
- Spreads centrifugally
- Red, sharply demarcated border
- Treatment: Topical antifungals
  - Clotrimazole, Miconazole
  - Terbinafine



Image courtesy of Robertgascoign/Wikipedia

## Tinea Versicolor

Pityriasis versicolor

- Cause by *Malassezia* species
- Dimorphic fungi, normal skin flora
- Yeast can transform to mycelial form → disease
- Transformation triggers:
  - Hot, humid weather
  - Sweating
  - Topical skin oils

## Tinea Versicolor


- Degradation of lipids → acids
- Damages melanocytes
- Hypopigmented skin
- Not a dermatophyte (does not consume keratin)



Image courtesy of Sarahrosenau/Wikipedia

## Tinea Versicolor

- KOH prep shows hyphae AND yeast cells
- “Spaghetti and meatballs”
- Treatment: Topical azoles
- Also, selenium sulfide (topical)
  - Promotes shedding stratum corneum



CDC (Dr. Lucille K. George)/Public Domain




Image courtesy of Dr. Lucille K. George /Wikipedia

## Sporothrix schenckii

Sporotrichosis

- Dimorphic yeast that lives on plants
- Spores introduced skin of hands with trauma
  - “Rose gardener’s disease”
- Papule at site of trauma days to weeks later
- Travels up arm via lymphatics
  - “Ascending lymphangitis”
- Similar lesions occur along lymph channels

## Sporothrix schenckii

Sporotrichosis

- Diagnosis: Fungal culture
- Treatment
  - Itraconazole (oral)
  - Saturated solution of potassium iodide (SSKI)

# Opportunistic Fungal Infections

Jason Ryan, MD, MPH

## Fungal Infections


- Pneumonia
  - Only in specific geographic areas
  - Histoplasmosis, Blastomycosis, Coccidioidomycosis
- Skin
  - Tinea versicolor, tinea pedis, sporothrix
- Opportunistic
  - Candida, Aspergillus, Cryptococcus, Mucormycosis, Pneumocystis

## Candida Albicans

- Normal flora of mouth, intestine, skin, vagina
  - Common contaminant of sputum culture
- Overgrowth disease
  - Oral thrush
  - Esophagitis
  - Vulvovaginitis
  - Diaper rash
- Disseminated disease
  - Endocarditis
  - Disseminated candidiasis


## Candida Albicans

- Dimorphic
- Forms pseudohyphae
  - Elongated, budding yeast cells
- Forms germ tubes ("germ tube test")
  - Yeast with hyphae growing out
  - Differentiates candida



Images courtesy of Y Tambe/Wikipedia

## Pseudohyphae



Images courtesy of Y Tambe/Wikipedia

## Candida

### Overgrowth Diseases

- Oral thrush
  - Inhaled steroid patients (asthma)
- Esophagitis
  - HIV/AIDS patients
  - White pseudomembrane on EGD
- Vulvovaginitis
  - "Yeast infection"
  - Itching, discharge ("cottage cheese" appearance)
  - Women taking antibiotics at risk (decreased normal flora)
- Diaper rash
  - Beefy, red plaques with satellite papules



James Heilman, MD/Wikipedia

## Candida

### Systemic Diseases

- Endocarditis (rare)
  - Almost always IV drug user
- Candidemia
  - From blood can spread to any organ system

## Candidemia

- Immunosuppressed patients
  - Neutropenic patients from chemo
- Patients in the ICU
- Central lines
- Total parenteral nutrition (TPN)/Hyperalimentation
- IV drug users

## Candida

### Treatment

- Vaginal disease/diaper rash
  - Topical azole
- Oral thrush
  - Nystatin "swish and swallow"
  - Fluconazole
- Esophagitis
  - Fluconazole
  - Resistant cases: voriconazole, caspofungin
- Candidemia/endocarditis
  - Fluconazole (stable, not immunocompromised)
  - Caspofungin or Amphotericin B

## Chronic mucocutaneous candidiasis

- Rare disorder
- Mutations in autoimmune regulator (AIRE) genes
- T-cell dysfunction
- T cells fail to react to candida antigens
- Chronic skin, mucous membrane candida infections
- Child with recurrent thrush, diaper rash

## Candida Immunity

- T-cells important for mucosal defense
  - Example: HIV patients often get thrush (LCD4)
- Neutrophils important for systemic defense
  - HIV patients rarely get candidemia
  - No candidemia in CMC
  - Chemo patients at risk for candidemia (neutropenia)

Pirofski L, Casadevall A; Rethinking T cell immunity in oropharyngeal candidiasis  
J. Exp. Med. Vol. 206 No. 2 269-273

## Aspergillus

- Aspergillus species ubiquitous in nature
  - A. fumigatus, A. flavus, and A. terreus
- Inhalation of spores (conidia) common
- Disease requires immunocompromise
- Usually chemo, stem cell transplant

## Aspergillus

- Catalase positive
  - Common infection in chronic granulomatous disease
- Monomorphic fungi
  - Do not form yeast cells
- Forms “branching septate hyphae”
  - V shaped branches
  - Visible septae
- Tips of some hyphae grow spores (conidiophore)

## Aspergillus




Image courtesy of Wikipedia/Public Domain



Image courtesy of KGH/Wikipedia




Image courtesy of CDC/Public Domain

## Aspergillus Disease

- Aspergillosis
  - Lung disease that can progress to systemic illness
  - Neutropenic patients
- Allergic bronchopulmonary aspergillosis
  - Allergic reaction in CF/Asthma patients
- Aspergilloma
  - Fungus invades pre-formed cavities (TB)
- Hepatocellular carcinoma
  - Aflatoxins

## Aspergillosis

- Severe lung disease
- Classic case:
  - Neutropenic patient
  - Fever, cough
  - Pleuritic chest pain
  - Hemoptysis
  - Multiple nodules/densities/infiltrates on imaging
- Can disseminate to any organ
  - Heart (endocarditis); Brain (abscesses; mycotic aneurysms)
- Treatment:
  - Voriconazole, Caspofungin, or Amphotericin B

## ABPA

Allergic bronchopulmonary aspergillosis

- Hypersensitivity (allergic) reaction to aspergillus
  - Type I (IgE)
  - Type III
- Lungs become colonized with Aspergillus
- Occurs predominantly in asthma and CF patients
- ABPA patients:
  - Increases Th2 CD4+ cells
  - Synthesis interleukins (IL-4, IL-5)
  - Eosinophilia
  - IgE antibody production

## ABPA

Allergic bronchopulmonary aspergillosis

- Classic case
  - Asthma or CF patient
  - Recurrent episodes cough, fever, malaise
  - Brownish mucus plugs, hemoptysis
  - Peripheral blood eosinophilia
  - High IgE level
- Diagnosis: Skin testing aspergillosis
- Treatment: Steroids

## Aspergilloma

- Fungus ball
- Caused by *Aspergillus fumigatus*
- Grows in pre-formed cavities
- Pulmonary TB is most common association
- Often asymptomatic
- Can cause hemoptysis
- Diagnosis: Imaging plus sputum culture
- Treatment: Observation vs. surgery

## Hepatocellular Carcinoma


- Aflatoxin produced by aspergillus
- Can contaminate corn, soybeans, and peanuts
- High rates of dietary intake associated with HCC
- Industrialized countries screen for aflatoxin
- Exposure from:
  - Food from non-industrialized countries
  - Locally grown foods

## Cryptococcus Neoformans

- Exists only as yeast
- Thick capsule
- Main disease is meningitis
  - HIV/AIDS
  - Immunocompromised (Chemo, post-transplant)
- Present in soil and pigeon droppings
- Inhaled → lungs → blood stream → meninges
- Rarely can cause pneumonia
- Rarely can spread to other tissues

## Cryptococcus Neoformans

- Can be cultured on Sabouraud's agar
- India ink staining shows capsules as "halos"
- Latex agglutination test
  - Detects polysaccharide capsular antigen



India Ink stain shows yeast with "halos"

Image courtesy of Crisco 1492

## Cryptococcal Meningitis

- Indolent symptoms over weeks
  - Fever, headache
- Can cause ↑ICP
- Risk of herniation with LP
- Must do CT or MRI
- Treatment:
  - Amphotericin B +/- Flucytosine
  - Fluconazole
- Sometimes intrathecal therapy used

## Mucormycosis

- Rare fungal infection of nose, eyes, brain
- Caused by *Rhizopus* sp. and *Mucor* sp.
- Fungi have enzyme: ketone reductase
- Thrive in high glucose, ketoacidosis conditions
- Serum from DKA patients stimulates growth
- Risk factors:
  - Diabetes, especially DKA
  - Treatment with steroids
  - Leukemia
  - Stem cell transplant patients

## Mucormycosis

- Disease from inhaled spores
- Enters nose or alveoli
- Angioinvasive fungus: Invades vessel walls
- Classically starts in sinuses
- Spreads to adjacent structures

## Mucormycosis

### Clinical features

- Severe sinusitis
  - Fever, discharge, congestion, sinus pain
- Necrosis of the palate
- Erythema/cyanosis of skin over sinuses
- Black eschars
- Orbital pain/swelling
- Facial numbness (cranial nerve damage)
- Cavernous sinus thrombosis

## Mucormycosis

### Clinical features

- Classic case:
  - Patient with DKA
  - Fever, headache, eye pain
- Diagnosis: mucosal biopsy
- Treatment:
  - Surgical debridement
  - Amphotericin B

## Mucormycosis

- Broad hyphae
- Irregularly branched, rare septations
- Different from *Aspergillus*:
  - Narrow, regular (v-shaped) branching, many septations




Image courtesy of Yale Rose/Flickr

## PCP


### *Pneumocystis jirovecii*

- Causes diffuse interstitial pneumonia
- Requires immunocompromise
  - Classically HIV
  - AIDS-defining illness
- Yeast → inhaled
  - Usually no symptoms if immune system intact
- CXR will show diffuse, bilateral interstitial infiltrates

## PCP

### *Pneumocystis jirovecii*

- Diagnosed by microscopy
  - Sputum sample, BAL, or biopsy
  - Sent for staining or fluorescent antibody testing
- Staining required to visualize → cannot be cultured
- Special stains used
  - Silver stains often used



CDC (Dr. Francis Chandler)/Public Domain




Image courtesy of Yale Rosen/Wikipedia

## PCP

*Pneumocystis jirovecii*

- Treatments
  - TMP-SMX (first line)
  - Dapsone
  - Pentamidine
- Prophylaxis
  - TMP-SMX when CD4 <200cells/microL
  - High dose steroid or other immunosuppressed patients

# Antifungal Drugs

Jason Ryan, MD, MPH

## Antifungal Drugs



- Amphotericin B
- Nystatin
- Flucytosine
- Azoles (fluconazole, itraconazole, voriconazole)
- Echinocandins (caspofungin, micafungin)
- Terbinafine
- Griseofulvin

## Antifungal Drugs


- Amphotericin drugs
  - Amphotericin B
  - Nystatin
  - Flucytosine
- Azoles (fluconazole, itraconazole, voriconazole)
- Echinocandins (caspofungin, micafungin)
- Tinea drugs
  - Terbinafine
  - Griseofulvin

## Sterols

- Steroids with alcohol groups present
- Cholesterol: animal cell walls
- Ergosterol: fungal cell walls



Ergosterol



Cholesterol

## Amphotericin B

- Binds ergosterol
- Forms pores in membrane
- Electrolyte leakage → cell death
- Used for dangerous, systemic fungal infections
  - Candidemia
  - Mucormycosis
  - Cryptococcus
  - Systemic histoplasmosis, blastomyces, coccidiomycosis
- Usually given intravenously
- Intrathecal administration meningitis

## Amphotericin B

- Several unique and important side effects
- Mechanisms not completely understood
- Many related to binding of cholesterol
- Fever, chills
  - "Shake and bake"
  - May be related to prostaglandin release
  - Minimize with Tylenol, NSAIDs, or diphenhydramine
- Phlebitis
  - Pain, inflammation of vein used for infusion
  - Avoided by using a central line
  - Sometimes hydrocortisone given with infusion
- Hypotension, arrhythmias

## Amphotericin B

- Nephrotoxicity
  - Causes renal vasoconstriction/toxic to tubules
  - Can insert into cell membranes → create pores
  - Decrease GFR (Cr will rise)
  - Rarely ARF
  - Hydration reduces this complication
- Liposomal Amphotericin B
  - Amphotericin B dissolved in lipids
  - Developed based on animal studies
  - Reduced incidence of nephrotoxicity

## Amphotericin B

- Hypomagnesemia, hypokalemia
  - Increased distal tubule permeability to Mg/K
  - Mg/K lost in urine
  - Need to replete Mg and K
- Anemia
  - Reversible, normocytic, normochromic anemia
  - Usually mild

## Amphotericin B

- Distal (Type I) RTA
  - Non anion gap metabolic acidosis
  - Very low HCO<sub>3</sub><sup>-</sup> (often <10meq/L)
  - Urine pH is high (pH>5.5)
- Nephrogenic diabetes insipidus
  - Hyponatremia


## Nystatin

- Binds ergosterol (same mechanism Ampho B)
- Highly toxic when given IV
- Not used systemically
- "Swish and swallow" for thrush (candida)
- Diaper rash (candida)

## Flucytosine

- Blocks fungal DNA/RNA synthesis
- Converted to 5-fluorouracil by cytosine deaminase
  - Cytosine deaminase only present in fungi
- Causes impaired DNA/RNA synthesis

## Flucytosine



## Flucytosine

- High incidence resistance when used alone
- Used in combination with Amphotericin B
- Main use is **cryptococcal meningitis**
- Major side effect is bone marrow suppression
  - Some spontaneous conversion to 5-FU
  - Leukopenia, thrombocytopenia

## Azoles

Fluconazole, itraconazole, voriconazole, clotrimazole, miconazole, ketoconazole

- Block ergosterol synthesis
- Inhibit P450 enzyme in fungi
- Enzyme converts lanosterol to ergosterol
- Side effects related to this mechanism:
  - Inhibits liver P450 system
  - Elevated levels of P450 meds (warfarin, theophylline)

## Azoles

- Hepatotoxicity
  - Reported with all azoles
  - LFTs monitored in patients on these drugs (oral)
- Ketoconazole
  - Life-threatening hepatotoxicity reported
  - Rarely used any longer for this reason
  - Suppresses cortisol synthesis (can be used in Cushing's)
  - Suppresses testosterone synthesis (causes gynecomastia)

## Azoles

- Itraconazole
  - Drug of choice for fungal pneumonias
  - Also Sporothrix
- Fluconazole
  - Excellent activity against Cryptococcus
  - Vulvovaginitis (Candida)
- Clotrimazole
  - Diaper rash
- Voriconazole
  - Severe, systemic fungal infections (Aspergillus)

## Echinocandins

Caspofungin, micafungin

- Inhibit cell wall synthesis
- Block synthesis of  $\beta$ -glucans (polysaccharides)
- $\beta$ -glucans account for 30-60% cell wall
- "Penicillin of antifungals"
- Used for severe, systemic infections
  - Aspergillus
  - Candidemia
- Few side effects
  - GI upset
  - Infusion-related histamine release (flushing)

## Terbinafine


- Blocks squalene epoxidase
- Key enzyme for ergosterol synthesis
- Given orally to treat dermatophyte infections
  - Especially onychomycosis
- Side effects:
  - Headache (13% patients)
  - Hepatotoxicity (monitor LFTs)
  - Rarely blurry vision

## Griseofulvin

- Blocks mitosis by interfering with **microtubules**
- Deposits in tissues with keratin
  - Binds to keratin → resistance to fungal invasion
- Oral therapy for skin/nail infections

## Griseofulvin

- Side effects:
  - Induces P450 (warfarin, theophylline levels will fall)
- Teratogenic: not safe in pregnancy
- Carcinogenic
- Other adverse effects:
  - Liver toxicity
  - Photosensitivity
  - Porphyria attacks



# Malaria

Jason Ryan, MD, MPH

## Malaria

- Protozoa infection of red blood cells and liver
- Occurs in tropics, subtropics
- Very rare in US, Europe
- Africa is most effected continent
- Transmitted by mosquito bite (female *Anopheles*)
- Caused by *Plasmodium* sp.
- Several species with distinct features:
  - *P. vivax*/*P. ovale*
  - *P. malariae*
  - *P. falciparum*

## Malaria

Life cycle

- Bite of female mosquito → sporozoites to liver
- Asymptomatic for up to 1 month

## Malaria

Life cycle

- Sporozoites invade hepatocytes
- Mature into multi-nucleated schizonts
  - "Pre-erythrocytic stage"

## Malaria

Life cycle

- Schizonts rupture → release Merozoites
- Invasion of RBCs ("Erythrocytic stage")

## Malaria

Life cycle

- Form trophozoites (ring form) in RBCs
- Inside RBCs mature to schizonts
  - Digest RBC proteins, especially hemoglobin
  - Breakdown products toxic to RBCs
- Merozoites formed (again) → RBC lysis
  - Occurs at regular intervals (48hr, 72hr)
  - Cyclic fevers can occur


## Malaria

### Life cycle

- Key Points:
  - Protozoa goes to liver after mosquito bite
  - Sporozoites are the infective form
  - Incubation period occurs
  - Release of merozoites leads to RBC infection, symptoms
  - Plasmodium matures/grows in RBC
  - Eventually ruptures RBC → release of merozoites
  - Cycle of maturation/release → cyclical fevers

## Malaria

### Life cycle



## Malaria

### Common symptoms

- Paroxysms of fever
  - Shivering and chills followed by high fever
  - Fever recurs at regular intervals (48hrs, 72hrs)
  - Variable by species of Plasmodium
- Anemia (RBC infection)
  - Severity varies by species of Plasmodium
  - Hemolytic: sometimes jaundice
- Splenomegaly
- Also nonspecific symptoms:
  - Sweating, fatigue, malaise, arthralgias, headache
  - Sometimes cough, vomiting, diarrhea

## Malaria

### Rare symptoms

- Altered consciousness (especially when febrile)
- Seizures
- "Blackwater fever"
  - Renal failure with hemoglobinuria
- Shock
- Severe symptoms usually due to *P. falciparum*

## *P. Vivax/Ovale*

- Classically has a 48hr cycle of fevers
  - "Tertian" fever pattern
  - Fever day 1, day 3
  - No fever day 2, day 4
- Dormant form in liver
  - "Hypnozoites" form
  - Recurring infection months after resolution
- Primaquine treats *P. vivax/ovale* liver disease
  - Without this, relapses may occur

## *P. malariae*

- Classically has a 72hr cycle of fevers
  - "Quartan" fever pattern
  - Fever day 1, day 4
  - No fever day 2, day 3

## P. falciparum

- Most severe malarial infection
- Fever pattern is irregular
- Invades RBCs of any age
  - Other forms invade only reticulocytes

## P. falciparum


- Induces sticky “knobs” on RBC surfaces
  - Knobs composed of parasite proteins
  - *P. falciparum* erythrocyte membrane protein 1 (PfEMP1)
- Knobs bind receptors on endothelial cells
- Result is occluded capillaries
  - Cerebral malaria (occluded vessels in brain)
    - Altered consciousness, delirium, coma
  - Renal failure (“blackwater fever”)

## Malaria


### Diagnosis

- Blood smear (Giemsa or Wright stains)

Trophozoite Ring



Schizont with merozoites



CDC/Public Domain

## Malaria Extras

- Duffy antigen
  - Necessary for *P. vivax* infection
  - Absence of Duffy → protective
- Sickle cell
  - May have evolved as protection from malaria
  - Children with HbS have lower risk of falciparum infection
- Thalassemia
  - Reduced parasite multiplication in *P. falciparum* infection

## Malaria

### Treatment

- Chloroquine
  - Weak base
  - Accumulates in food vacuoles (acidic) of RBC trophozoites
  - Blocks plasmodium **heme polymerase**
  - Heme portion of Hgb toxic to parasite
  - Plasmodium converts this to nontoxic form
- Lots of chloroquine resistance
- Used mainly in limited areas (“chloroquine sensitive”)
- Not used for severe infections
- Only kills erythrocytic forms (not liver forms)
- Retinopathy associated with long-term use

## Malaria

### Treatment

- Severe infections
  - Artesunate (IV)
  - Quinidine (IV)
- Other drugs
  - Mefloquine (commonly used in chloroquine resistant areas)
  - Primaquine (liver phase only; not active against RBC phase)
  - Atovaquone

## Immune Suppression

- Chloroquine and hydroxychloroquine
- Malaria drugs with immunosuppressive actions
  - Block TLRs in B-cells (↓activation)
  - Weak bases: ↑pH in immune cells → ↓ activity
  - Other actions
- Used in rheumatoid arthritis, SLE

## G6PD Deficiency

- X-linked genetic disorder
- Hemolytic anemia triggered by various stressors
  - Infections
  - Fava beans
  - Drugs
- Many malaria drugs trigger anemia in G6PD
  - Quinidine
  - Primaquine
- Often test for G6PD deficiency prior to treatment:
  - Primaquine for *P. vivax/ovale* liver phase
  - IV Quinidine for life threatening *P. Falciparum*

# Protozoa

Jason Ryan, MD, MPH

# Protozoa

- Protozoa are eukaryotes
  - Have a nucleus
  - Intracellular organelles
  - VERY different from bacteria
- Unicellular
- Mobile
- Easily seen under microscope

# Protozoa

- Exist in different stages
- Trophozoites
  - Feeding form
  - Vulnerable to environmental conditions
- Cysts
  - More durable form
  - Often present in feces → water → new infection

# Protozoa Infections

- GI Illness
  - Giardia, Entamoeba, Cryptosporidium
- CNS Infections
  - Toxoplasma, Naegleria fowleri, Sleeping sickness
- Blood infections
  - Malaria, Babesia
- Others
  - Chagas disease, Leishmaniasis, Trichomonas

# Protozoa

## Transmission

- All GI protozoa transmit fecal → oral
  - Cysts in stool → water
  - Consumption of contaminated water
- Others transmitted by various methods
  - Direct (Trichomonas; STD)
  - Contact with cat feces (Toxoplasmosis)
  - Mosquito/fly (Malaria, Babesia)

# Protozoa Drugs

- Metronidazole works for many infections
  - GI parasites: Giardia, Entamoeba
  - Trichomonas
- Most other drugs unique to one protozoa
  - Iodoquinol (Entamoeba)
  - Nitazoxanide (Cryptosporidium)
  - Suramin (Trypanosomes)
  - Melarsoprol (Trypanosomes)
  - Atovaquone (Babesia)
  - Nifurtimox (Chagas disease)
  - Sodium stibogluconate (Leishmania)

## Giardia Lamblia

### Giardiasis

- Cysts found in moist environments
- Classic source is water from a mountain stream
- Ingested cysts → trophozoite in intestine
- Affects small intestine
- Bloating, foul smelling, fatty diarrhea
- Steatorrhea
- Stools that float

## Giardia

- Diagnosis:
  - Cysts in stool
  - Trophozoites in stool
  - ELISA for Giardia antigens in stool
- Classic case: Camper/hiker, diarrhea, flatulence
- Treatment: Metronidazole



Image courtesy of Joel Mills/Wikipedia



Image courtesy of Doc. RNDr. Josef Reischig, CSC/Wikipedia

## IgA Deficiency

- IgA very important for defense against Giardia
- Lack of IgA → Recurrent/chronic giardia infection
- Bruton's Agammaglobulinemia
- Selective IgA deficiency

Langford TD et al. Central Importance of Immunoglobulin A in Host Defense against Giardia spp. Infect. Immun. January 2002;vol. 70 no. 1 11-18

## Entamoeba Histolytica


### Amebiasis/Amebic dysentery

- Found worldwide
- Common in developing countries/poor sanitation
- Cysts ingested in contaminated water
- Form trophozoites in small intestine and invade tissue
- Causes bloody diarrhea (dysentery)
- Ascends portal system → liver
- Liver abscesses
  - RUQ pain
  - "Anchovy paste" exudate
- Traveler, bloody diarrhea, RUQ pain → Entamoeba

## Entamoeba Histolytica

- Diagnosis:
  - Stool microscopy
  - Serology (antibodies to Entamoeba)
- Treatment:
  - Metronidazole
  - Iodoquinol (asymptomatic cysts carriers)

Trophozoite with ingested RBCs



Wikipedia/Public Domain


Cysts with multiple nuclei



Wikipedia/Public Domain

## Entamoeba Histolytica

"Flask-shaped" ulcer on biopsy



CC0[Dr. Mae Mevin]/Public Domain

## Entamoeba Histolytica


- Classic case
  - Patient in a developing country (or recent immigrant)
  - Bloody diarrhea developing over weeks
  - RUQ pain

## Cryptosporidium

- Protozoa forms eggs (oocysts)
- Found in contaminated water
- Ingestion → infection
- Chlorination does not destroy oocysts
- Infection in swimming pools common
- Immunocompetent patients
  - Mild, watery diarrhea
  - Self-limited
- HIV/AIDS
  - Severe diarrhea


## Cryptosporidium

- Diagnosis: Microscopy
  - Acid fast staining reveals oocysts
- Treatment:
  - Nitazoxanide (only in immunocompetent)
  - Anti-retroviral therapy for HIV patients
- Prevention is key
  - Wash hands
  - Filter water



Wikipedia/Public Domain

## GI Protozoa



## Toxoplasma gondii

Toxoplasmosis

- Commonly lives in cats (felines)
- Oocysts shed in stool
- Infection from ingested oocysts (soil)
- Also meat from contaminated animal (cysts)
- Invades intestine → disseminates
- May enter latent phase → reactivate later
- Two major disease processes
  - HIV CNS disease
  - Congenital toxoplasmosis

## Toxoplasma gondii

Toxoplasmosis

- Significant CNS disease immunosuppressed
  - Usually HIV/AIDS (CD4 <100cells/mm<sup>3</sup>)
  - Sometimes "reactivates"
- Brain abscesses (fever, headache, nerve palsies)
- Multiple "ring-enhancing" lesions on imaging




Image courtesy of LearningRadiology.com

## Toxoplasma gondii

### Toxoplasmosis


- Crosses the placenta
- Range of symptoms/signs in fetus
- Classic triad:
  - Chorioretinitis (inflammation of choroid in eye)
  - Hydrocephalus
  - Intracranial calcifications (seen on imaging)

## Toxoplasma gondii

### Diagnosis

- Serology
  - IgG or IgM antibodies to Toxoplasma
  - IgM antibodies appear within one week, rise, decline
  - IgG antibodies rise within two weeks, generally persist for life
- Biopsy

Tachyzoite (Giemsa stain)



CDC/Public Domain

## Toxoplasma gondii

### Toxoplasmosis

- Treatment:
  - Sulfadiazine/pyrimethamine
- Blocks THF synthesis pathway
- Similar to TMP/SMX

## Naegleria fowleri

- Rare cause of fatal meningoencephalitis
  - 300 cases reported worldwide
- Found in freshwater lakes/ponds
- Contaminated water → nose → cribriform plate
- Classic case
  - Recent (4-5 days ago) swimming
  - Fever, confusion, stiff neck
  - Often fatal (99% in one series)

## African trypanosomiasis

### Trypanosoma brucei, T. gambiense, T. rhodesiense

- Protozoa infections from insect bite
- All occur in Africa
- All caused by tsetse fly
- "African sleeping sickness"
- Early and late features
  - Early: fever, arthralgias
  - Late: Somnolence, coma
- Organisms visible on blood smears



CDC/Public Domain

## African trypanosomiasis

### Trypanosoma brucei, T. gambiense, T. rhodesiense

- Key feature: recurring fever
- Due to antigenic variation
- "Variant surface glycoproteins" (VSG)
  - Each trypanosome covered ~10million copies of one VSG
- Change VSG when host mounts immune response
- Waves of parasitemia
- Recurring fever

## Babesia

### Babesiosis

- Transmitted by Ixodes tick
- Same tick that transmits:
  - Borrelia (Lyme)
  - Anaplasma (Anaplasmosis)
  - Co-infection common
- Same geography as Lyme: Northeastern US
- Infects red blood cells
- Increased risk in asplenic patients
  - Spleen clears Babesia/infected RBCs

## Babesia

### Babesiosis

- Fever
- Hemolytic anemia
- Splenomegaly


## Babesia

### Babesiosis




Maltese Cross

- **Diagnosis:**
  - Blood smear (ring forms; Maltese crosses)
  - PCR (amplification babesia RNA)
- **Treatment:**
  - Azithromycin (macrolide)
  - Atovaquone (malaria drug)



CDC/Public Domain



CDC/Public Domain

## Trypanosoma cruzi

### Chagas' disease

- Transmitted by reduviid bug
- Found in South America
- Bugs nest in cracks/holes of housing
- Acute phase – nonspecific, febrile illness
- Chronic Chagas: heart, esophagus, colon



CDC/Public Domain

## Trypanosoma cruzi


### Chronic Chagas' disease

- **Cardiac**
  - Right and left heart failure
  - High prevalence ventricular thrombi
  - Pulmonary embolism/stroke
- **Esophagus**
  - Achalasia, megaesophagus (dilation)
- **Colon**
  - Megacolon (severe constipation)

## Trypanosoma cruzi

### Chagas' disease

- **Acute phase: blood smear**
  - Trypomastigotes visible
- **Chronic phase**
  - Serology (IgG antibodies)
- **Treatment: Nifurtimox**
  - Acute phase
  - Not effective with advanced disease



CDC/Public Domain

## Leishmania donovani

### Leishmaniasis

- Transmitted by sand fly
- Mostly Asia, Africa, South and Central America
- Protozoa infects macrophages
- Cutaneous leishmaniasis
  - Large ulcer with indurated borders
- Visceral leishmaniasis
  - Kala-azar (Hindi: "black fever")
  - Fever
  - Painful splenomegaly
  - Pancytopenia




CDC/Public Domain

## Leishmania donovani

### Leishmaniasis

- Diagnosis by biopsy of affected organs
  - Usually bone marrow or spleen
- Amastigotes in macrophages
  - Small, round or oval bodies
- Treatment:
  - Amphotericin B
  - Sodium stibogluconate




CDC/Public Domain

## Trichomonas vaginalis

- Lives in urogenital tract
- Sexually transmitted (no cyst form)
- Men:
  - Usually asymptomatic
  - Can cause urethritis (discharge, dysuria)
- Women
  - About 50% asymptomatic
  - Vaginitis
  - Itching
  - Classically yellow-green, foul-smelling discharge

## Trichomonas vaginalis

- Diagnosis:
  - Wet mount: motile trichomonads
  - pH >4.5 (normal 4-4.5)
- Treatment: Metronidazole
  - Patient and partner
- One of 3 main causes vaginitis
  - Bacterial vaginosis (*Gardnerella vaginalis*)
  - Candida (fungi)
  - Trichomonas (protozoa)



CDC/Public Domain



Clue Cell - Bacterial Vaginosis

CDC/Public Domain

## Vaginitis

	Bacterial vaginosis	Candida	Trichomonas
Discharge	Thin, off-white	Thick, White "Cottage Cheese"	Yellow-green
Odor	Fishy	None	Foul-smelling
Tissue	Normal	Vaginal erythema	Strawberry cervix
pH	>4.5	Normal (4-4.5)	>4.5
Other test	Whiff test	KOH Prep	Wet Mount
Treatment	Metronidazole	Fluconazole	Metronidazole

Whiff test: KOH yields fishy odor  
 KOH Prep: Shows pseudohyphae in candida  
 Wet mount: Motile trichomonads


# Helminths

Jason Ryan, MD, MPH

## Helminths

Worms

- Roundworms (nematodes)
- Flatworms
  - Tapeworms (cestodes): Ribbon-like
  - Flukes (trematodes): Look like leaves
- All have three stages
  - Eggs
  - Larvae
  - Adults



## Helminths

- Very rare in developed world
- Most cause eosinophilia
- Various modes of infection:
  - Ingestion of eggs
  - Penetration of skin

## Helminths

- Many unique drugs used for therapy
- Bendazoles
  - Albendazole
  - Mebendazole
- Others:
  - Ivermectin
  - Pyrantel pamoate
  - Diethylcarbamazine
  - Praziquantel

## Helminths

Things to know

- Name of organism
- Symptoms
- Mode of infection
- Diagnosis (often stool analysis)
- Treatment

## Enterobius vermicularis

Pinworm

- Most common helminth infection US
- Common among children
- Eggs found in moist environments
- Child touches eggs, contaminates fingers
- Fingers touch food, mouth → ingestion of eggs
- Eggs hatch in small intestine
- Adults deposit eggs in perianal folds

## Enterobius vermicularis

Pinworm

- Most infections asymptomatic
- Most common symptom: perianal itching
  - Inflammatory reaction to worms and eggs on skin
  - Occurs predominantly at night
- Diagnosis: Scotch tape test
  - Adhesive applied to perianal skin
  - Placed on glass slide
  - Eggs visualized under microscope
- Treatment:
  - Bendazoles (albendazole, mebendazole)
  - Pyrantel pamoate

Children, itchy anus, Scotch tape test

## Ascaris lumbricoides

Giant roundworm

- Found in warm, tropical climates
- Common in children (vomiting worms!)
- Worms live in small intestine of infected patients
- Shed eggs in stool
- Eggs survive in environment
  - Fecal-oral transmission
- When ingested eggs hatch in small intestine
- Release larvae → penetrate intestinal wall
- Migrate via blood or lymphatics

## Ascaris lumbricoides

Giant roundworm

Ascaris Egg



CDC/Public Domain

- Most patients asymptomatic
- Intestinal symptoms
  - GI upset
  - Bowel obstruction
- Pulmonary symptoms
  - Loeffler's syndrome
  - Eosinophilic pneumonitis from worm migration to lungs
- Diagnosis: Eggs seen on stool examination
- Treatment:
  - Bendazoles (albendazole, mebendazole)
  - Pyrantel pamoate

## Ascaris lumbricoides

Giant roundworm

- Classic case
  - Patient with recent travel
  - Abdominal pain
  - Wheezing, cough
  - Eosinophilia
  - Eggs seen on stool examination

Bowel obstruction, pneumonia, eggs in stool

## Strongyloides stercoralis

- Larvae found in soil
- Penetrate skin
- Migrate via blood to lungs
  - Penetrate alveolar air sacs
- Ascend tracheobronchial tree → swallowed
- Mature into adults, burrow into duodenum/jejunum

## Strongyloides stercoralis

- Skin reactions
  - Rash, often severe itching
- Pulmonary migration
  - Dry cough, throat irritation, dyspnea, wheezing, hemoptysis
- Duodenitis
  - Upper abdominal pain, diarrhea, anorexia, nausea, vomiting
- Diagnosis: Stool larvae or serology
- Treatment:
  - Albendazole
  - Ivermectin

Skin, belly pain, cough

## Hookworms

*Ancylostoma duodenale*, *Necator americanus*

- Worms live in small intestine of infected patients
- Shed eggs in stool
- Eggs hatch in soil → larvae
- Larvae penetrates skin
- Migrate into blood, carried to lungs
- Ascend bronchial tree → swallowed
- Mature to adults in intestine

## Hookworms

*Ancylostoma duodenale*, *Necator americanus*

- Major impact is on nutritional status
- Worms attached to intestinal mucosa
- Cause blood loss by ingesting blood
  - Facilitated by production of anticoagulants
- Daily losses of blood, iron, and albumin
- Result: Anemia, malnutrition
- Diagnosis: Stool exam for eggs
- Treatment:
  - Benzodoles (albendazole, mebendazole)
  - Pyrantel pamoate

Skin, skinny, anemia

## Trichinella

Trichinosis



- Cysts in undercooked meat
- Larvae invade small bowel → adults
- Migrate to striated muscles
- Symptoms: muscle weakness
- Diagnosis: serology, biopsy
- Treatment: Benzodoles

Meat, muscles

## Intestinal Nematodes

Summary

- Ingested eggs: Enterobius, Ascaris, Trichinella (cysts)
- Skin penetration: Strongyloides, Hookworms
- Most diagnosed with stool examination
  - Exception: Pinworm (tape test)
- Most treated with benzodoles and pyrantel pamoate
  - Exception: Strongyloides (Ivermectin/albendazole)

## Dracunculus medinensis

Guinea worm

- Consumption of unfiltered water
- Water contains copepods (small crustaceans)
- Copepod dies, spills larvae into intestine
- Female adults migrate to skin
- Up to a year later, worm migrates to surface of skin
- Painful papule develops
- Worm emerges → burning sensation
- Treatment: Extraction of worm slowly
  - Can take days or weeks!

Giant skin worm

## Dracunculus medinensis



CDC/Public Domain

## Onchocerca volvulus

River blindness

- Infection from female blackfly bite
- Deposits larvae into skin
- Mature into adults
- Adults produce offspring (microfilariae)
- Microfilariae move through tissues
  - Subcutaneous, dermal, ocular, lymph system
- Provoke a mild immune response while alive
- When they die, significant inflammatory response

## Onchocerca volvulus

River blindness

- Skin symptoms
  - Generalized itching
  - Subcutaneous nodules ("onchocercoma")
  - Many other skin symptoms possible
- Eye
  - Keratitis, uveitis, blindness
- Diagnosis:
  - Skin biopsy ("skin snips")
  - Examined for microfilariae
- Treatment: Ivermectin

Itching, blindness, skin snips

## Loa Loa

Loiasis

- African eye worm
- Transmitted by biting deerflies (horse fly or deer fly)
- Fly introduces larvae to skin
- Larvae penetrate bite wound
- Mature into adult worms over months
- Adults live in the subcutaneous tissue
- Migrate to other areas, especially eye

## Loa Loa

Loiasis



Lichtinger A, Caraza M, Halpert M - Am. J. Trop. Med. Hyg. (2011)  
Open Access - Creative Commons

## Loa Loa

Loiasis

- Most individuals asymptomatic
- Two main clinical manifestations:
  - Subcutaneous swellings (Calabar swellings)
  - Migration of worms across subconjunctiva of eye
- Worms measure 3 to 7 cm
- Can be visualized directly crossing the conjunctiva
  - Often takes 10 to 20 minutes!
- Diagnosis:
  - Visualizing adult worm: subcutaneous tissue or conjunctiva
  - Blood smear: Detection of microfilariae
- Treatment: Diethylcarbamazine

Eye worm, skin swelling

## Lymphatic Filariasis

*Wuchereria bancrofti*, *Brugia malayi*, *Brugia timori*

- Transmitted by mosquito bites
- Larvae migrate to lymphatic system
- Grow into adults over months (up to 1 year)
- Obstruct of lymphatic flow

## Lymphatic Filariasis

*Wuchereria bancrofti*, *Brugia malayi*, *Brugia timori*

- Lymphedema/Elephantitis
  - Massive non-pitting edema
  - Hardening of tissues
  - Hyperpigmentation
- Major cause of disfigurement/disability
- Diagnosis: microfilariae seen on blood smear
- Key finding is eosinophilia
- Treatment: Diethylcarbamazine



Elephantitis

## Toxocara

Visceral larva migrans

- Not natural human parasites
  - *Toxocara canis* → dogs
  - *Toxocara cati* occurs in cats
- Disease of young children
  - Exposed to playgrounds/sandboxes contaminated by pet feces
- Hepatitis and pneumonitis
  - Larvae migrate to liver and lungs
- Key finding: eosinophilia
- Diagnosis: Serology
- Treatment: Bendazoles (albendazole, mebendazole)

## Toxocara

Visceral larva migrans

- Classic case
  - Child who plays in sandbox, eats dirt
  - Mention of cat or dog exposure
  - Wheezing, dyspnea (often no history of asthma)
  - RUQ pain, hepatomegaly

Cats, dogs, kids, liver, lungs

## Taenia solium

Taeniasis


- Larval form (oncosphere) in raw, undercooked meat
- Ingested → matures into intestinal tapeworm
- Most infections asymptomatic
- Sometimes mild intestinal symptoms
  - Nausea, anorexia, epigastric pain
- Eosinophilia occurs
- Diagnosis:
  - Eggs or worms segments (proglottids) in stool
- Treatment: Praziquantel

Meat tapeworm in gut

## Cysticercosis

- Caused by **eggs** (cysts) of *Taenia solium*
- Not from undercooked pork
- Tapeworm carriers can shed eggs in stool
- Fecal-oral ingestion of eggs that hatch in intestine
  - Often another member in household with carrier
- Invade the bowel wall and disseminate
  - Brain, muscles, liver
- Over weeks, tissue forms (cysticerci) develop
  - Membranous walls filled with fluid
- Cysts in brain → neurocysticercosis
- Major cause of seizures in underdeveloped countries

## Cysticercosis



CDC/Public Domain

Eggs, brain cysts, seizures

## Diphyllobothrium latum

- Tapeworm similar to *Taenia solium*
- Transmission from eating infected fish
- Classic unique feature: Anemia
  - *D. latum* has affinity for vitamin B12
  - Competes with host for vitamin
- Macrocytic, megaloblastic anemia
- Hypersegmented PMNs
- Neurologic symptoms
  - Paresthesias, subacute combined degeneration

B12 deficiency

## Echinococcus granulosus

- Dogs are definitive host
- Eggs shed in dog stool
- Fecal-oral ingestion of eggs
- Eggs hatch, penetrate intestinal mucosa
- Enter blood/lymphatic system

## Echinococcus granulosus

- Main clinical problem is liver cysts
- Can become massive (>10cm!)
- Hepatomegaly, RUQ pain
- Cysts may rupture → fever
  - May cause acute hypersensitivity reactions
  - Sometimes anaphylaxis
- Treatment: surgery
  - Pre-inject with agent to kill parasite
  - Hypertonic saline, ethanol
  - Adjunctive therapy with albendazole

Giant liver cysts

## Schistosoma

Schistosomiasis

- Worms live in snails
  - Infectious form (cercariae)
  - Emerge from the snail, contaminate water
  - Cercariae penetrate skin of humans
- Over weeks, worms migrate through tissue and develop into adult worms inside blood vessels
- Mature worms produce eggs in bladder, intestine, spleen, liver
- Inflammation/scarring over years

## Schistosoma

Schistosomiasis

- Acute infection
  - Swimmer's itch
  - Hypersensitivity (fever, urticaria and angioedema)

## Schistosoma

### Schistosomiasis

- Chronic infection: multisystem
  - GI, liver, spleen, GU, lungs, CNS
- GI
  - Abdominal pain, blood in stool
- GU:
  - Hematuria
  - Squamous cell carcinoma
- Liver/Spleen
  - Hepatosplenomegaly, Portal hypertension
- Granulomas
- Treatment: Praziquantel

Snails, skin, squamous cell, granulomas

## Clonorchis sinensis

### Chinese liver fluke

- Korea, Japan, Taiwan, and Southern China
- Infection from eating contaminated fish
  - Eggs hatch in snails, develop in cercariae
  - Cercariae released from snails to water
  - Penetrate flesh of fish
  - Humans eat fish → illness
- Flukes ascend and reside in biliary tract

## Clonorchis sinensis

### Chinese liver fluke

- Biliary tract inflammation and obstruction
  - Obstructive jaundice
  - Pancreatitis
- Two special complications:
  - Pigmented (bilirubin) gallstones
  - Cholangiocarcinoma
- Praziquantel

Fish in the gall bladder

## Paragonimus westermani

- Raw or undercooked crayfish or crabs
- Fluke migrates to lungs
- Recurrent hemoptysis
  - Chocolate colored sputum (blood, inflammatory cells, eggs)
- Secondary bacterial infections common
- Diagnosis: Eggs in sputum or lavage
- Treatment: Praziquantel

Coughing crabs

# Virus Structure

Jason Ryan, MD, MPH

## Viruses

- Nucleic acids (either DNA or RNA)
- Surrounded by protein called a capsid
- Sometimes surrounded by envelope
- No metabolic activity

## Virus Structure




Image courtesy of Emmanuel Boutet/Wikipedia

## Capsids

- Most capsids have one of two common shapes




Image by Dr. Richard Feldmann, Wikipedia/Public Domain




Image courtesy of Arionfx/Wikipedia

## Envelopes

- Viruses are “naked” or “enveloped”
- Envelope: lipid membrane acquired from host cell during assembly of the virus
  - Usually host cell plasma membrane
  - Sometimes nuclear membrane, endoplasmic reticulum
- Viral glycoproteins often embedded in membrane
  - Used for binding to host cells
  - Also antigens for immune system
- All naked viruses have icosahedral capsids
- Enveloped have icosahedral or helical

## Envelopes

- | Enveloped     | Naked           |
|---------------|-----------------|
| • Hepatitis B | • Adenovirus    |
| • Herpes      | • Rotavirus     |
| • HIV         | • Rhinovirus    |
|               | • Hepatitis A/E |


## RNA Virus Genomes

- Most are single stranded and linear
  - Exception: Reoviruses (ds RNA)
  - Exception: Retroviruses (single stands x 2)
  - Circular: Bunyaviruses, arenaviruses, delta virus (BAD)
- Most replicate in cytoplasm
  - Exception: Influenza, Retroviruses
  - Replicate in nucleus

## RNA Virus Polarity

- Can be (+) sense or (-) sense (i.e. polarity)
- Positive stranded RNA
  - Structurally similar to mRNA
  - In cytoplasm, used for protein synthesis immediately
- Negative stranded RNA
  - Must be converted to (+) RNA first
  - Can then be used as template for protein
  - Virus must carry enzyme to convert (-) to (+) RNA


## RNA Orientation



RNA Bases: Uracil, Guanine, Adenine, Cytosine

Narayanan/Wikipedia


## RNA Virus Polarity




## RNA Polymerase

- Human cells make RNA from DNA
  - Transcription
  - Enzyme: "DNA-dependent RNA polymerase"
- Viruses make RNA from RNA
  - Must synthesize their own enzyme
  - "RNA-dependent RNA polymerase"

## RNA Virus Replication




## RNA Virus Replication



## Retroviruses (RNA)

HIV, HTLV



## DNA Virus Genomes

- Circular or linear DNA
- Most have double stranded DNA
  - Except **parvovirus** which is single stranded


## Viral Genome Infectivity

- (+) RNA genomes: Infectious by themselves
  - Genetic material begins producing new proteins/nucleic acids on entry into cell
- dsDNA genomes: Infectious by themselves
- (-) RNA: NOT infectious by themselves
  - Require RNA-dependent RNA polymerase to reproduce


## Segmented Genomes

- Many RNA viruses are segmented
- Multiple molecules of RNA ("segments") in virus
- Allows for re-assortment of RNA
  - Two viruses co-infect same cell
  - Mixing of segments into new virus
- Most important example is influenza virus
- BOAR
  - Bunyaviruses (California, Congo, Hanta)
  - Orthomyxovirus (Influenza)
  - Arenavirus (LCMV, Lassa)
  - Reovirus (Rotavirus, Coltivirus)

## DNA Viruses



All DNA viruses (+) have icosahedral capsid except Pox  
All DNA viruses replicate in nucleus except Pox




- ## Viral Vaccines
- **Live, attenuated vaccines**
    - Inactive strains
    - Rarely can produce clinical disease
    - Cannot give to immunocompromised
    - Long lasting protection (no boosters)
    - Smallpox, yellow fever, chickenpox (VZV)
    - Sabin's polio virus, MMR, Intranasal influenza
  - **Killed virus vaccines**
    - No risk of infection
    - Less immune response (boosters often required)
    - Rabies, Injected influenza, Salk Polio, HAV

- ## Viral Vaccines
- **Recombinant vaccines**
    - Vaccine protein gene inserted into a virus or cells in culture
    - When carrier virus or cell grows, vaccine protein created
    - Immune system will recognize the expressed protein
    - Hepatitis B vaccine: Recombinant HbsAg
    - HPV: Recombinant proteins types 6, 11, 16, and 18

# DNA Viruses

Jason Ryan, MD, MPH

## DNA Viruses



All DNA viruses (+) have icosahedral capsid except Pox  
 All DNA viruses replicate in nucleus except Pox All  
 DNA viruses are ds except Parvovirus (ss) All DNA  
 viruses are linear except middle 3

## Parvovirus

- Very small virus
- Non-enveloped
- Single-stranded (-) DNA virus (only one!)
- B19 is predominant parvovirus in humans
- Four important syndromes
  - Aplastic crisis in sickle cell anemia
  - Hydrops fetalis
  - Fifth disease in children
  - Arthritis in adults

## Aplastic Crisis

- B19 replicates in RBC progenitor cells
  - Only replicates in S phase (no S phase in mature RBCs)
  - Causes "S phase arrest"
- Bone marrow and blood → ↓erythropoiesis
- Healthy patients:
  - RBC production returns 10 to 14 days; mild/no anemia
- Sickle cell patients
  - Increased RBC turnover
  - Lack of erythropoiesis leads to severe anemia
  - Pallor, weakness, and lethargy

## Aplastic Crisis

- Watch for a sickle cell patient with LOW retic count
  - Normal reticulocyte count 0.5 to 1.5%
  - Should be high in anemia
  - If low, think B19
- Symptoms of anemia: fatigue, dyspnea
- Treatment: Transfusions
- Infection resolves days → weeks

## B19 in Pregnancy

- Fetus especially vulnerable to B19
  - Shortened RBC half-life
  - Expanding RBC volume
  - Immature immune system
- B19 infection in pregnancy: miscarriage, fetal death
- Hydrops fetalis
  - Fluid accumulation in fetus (ascites, pleural, etc.)
  - Diagnosed on ultrasound
  - "Immune hydrops" from Rh mismatch
  - Many non-immune causes including B19

## Fifth Disease

Erythema infectiosum; slapped cheek disease

- Mild fever, rash in children
- Outbreaks among school aged children
- Fever, runny nose (due to viremia)
- Followed by rash (few days later)
  - Probably immune related
  - Viral infection has usually resolved
- Cheeks look like they have been slapped
- Face rash often followed later by rash on trunk/limbs
- No diagnostic test or treatment (self limited)
- Adults may catch this: mild arthralgia/arthrits

## Arthritis

- B19 can cause acute arthritis
- Often in adults, usually women
- About 75% will have rash
  - Various rashes
  - Usually not slapped cheeks
- Symmetric, most frequently in small joints
  - Hands, wrists, knees, feet
- Joint stiffness is common (can mimic RA)
- Diagnosis: B19 antibodies in plasma
- Usually resolves in few weeks

## Adenovirus

- Non-enveloped, icosahedral
- Double stranded DNA virus
- Important syndromes
  - Pharyngitis, Pneumonia
  - Pink eye (conjunctivitis)
  - Hemorrhagic cystitis
- Very stable - survive on surfaces
- Transmission:
  - Aerosol droplets
  - Fecal-oral
  - Contact with contaminated surfaces



Image courtesy of Joyhill09/Wikipedia

## Acute Hemorrhagic Cystitis

- Cause by adenovirus
- Occurs in children
  - Watch for outbreaks at day care centers/schools
- Hematuria, sometimes gross
- Sometimes dysuria
- Usually no fever, other symptoms
- Self-limited

## Papillomavirus

- Non-enveloped
- Double stranded, circular DNA virus
- Multiple subtypes: 1, 2, 6, 11, 16, 18
- Clinical disease (depends on subtype):
  - Cutaneous warts
  - Genital warts
  - Cancer

## Cutaneous Warts

- Caused by papillomavirus (1, 2, 3, 4, 7, 10)
- Treatment: salicylic acid or liquid nitrogen



Steven Fruitsmaak/Wikipedia

## Anogenital Warts

Condylomata acuminata

- STD caused by papillomavirus (6, 11)
- Soft, tan, cauliflower-like lesions
- "Verrucous" = warts
- Penis, vulva, perianal area (rectal bleeding)
- Treatment:
  - Chemical agents
  - Surgical therapy
- Does not lead to cancer



SOA-AIDS Amsterdam/Wikipedia

## HPV and Cancer


- Persistent infection over years can lead to cancer
- Malignancies associated with HPV infection:
  - Cervical
  - Anal, Penile
  - Oropharyngeal squamous cell cancers (mouth, throat)
- Usually types 16 and 18
  - Responsible for about 70 percent cases
- All more common in HIV/AIDS

## Cervical Cancer

- High prevalence HPV among sexually active women
  - Most will clear infection
  - Some will have infection persist
- Vaccine available (capsid proteins)
  - Some target types 16/18
  - Others also target 11/6 (genital warts)
- Screening done with Pap smear

## Koilocytes

- Seen on Pap smear
- Epithelial cell infected by HPV
- Large, darkened nuclei
- Perinuclear haloes



Public domain/Wikipedia


## Polyomavirus

- Non-enveloped
- Double stranded DNA virus
- Circular
- Disease in immunocompromised patients
- JC Virus: PML
  - Progressive multifocal leukoencephalopathy
  - CNS disease in HIV patients
- BK Virus
  - Classic disease in post-kidney transplant patients
  - Slowly progressive rise in creatinine

## Hepadnavirus

Hepatitis B virus

- Enveloped, Circular
- #1: Partially double stranded DNA virus
  - Genome enters hepatocytes → nucleus
  - DNA becomes fully double stranded
  - mRNA synthesized → cytoplasm
- #2: Reverse transcriptase synthesized
  - Viral mRNA → viral DNA
  - Packaged in capsid
- #3: Envelope from endoplasmic reticulum



## Poxvirus

Dr. Graham Beards/Wikipedia

- Enveloped
- Double stranded DNA virus
- Linear
- Capsid not icosahedral or helical
  - Complex shape: either an oval or brick-shape
- Large virus
- Replicates in cytoplasm (not nucleus!)
- Virus contains DNA-dependent RNA polymerase
- Makes RNA in cytoplasm
- Synthesis of proteins for replication

## Poxvirus

- Relevant diseases:
  - Smallpox
  - Molluscum contagiosum

## Smallpox

Variola virus

- Initially fevers, headache, malaise
- Skin rash erupts, goes through phases
- Macules → papules → raised pustules
- Eradicated by vaccination 1970s
- Concern for bioterrorism
  - Virus maintained US/Russian labs
  - Concern for release
  - Possible hidden stockpiles

CDC/Public Domain

## Cowpox

Cowpox virus

- Causes pustules on cows
- Milkmaids often got small blisters on their hands
- Edward Jenner inoculated a boy with cowpox (1796)
- Then exposed him to smallpox → no infection

## Vaccinia

Pox Virus

- DNA virus in the pox family
- Causes mild skin reaction
- Used to vaccinate against smallpox

## Molluscum Contagiosum

Molluscum Contagiosum Virus

- Member of poxvirus family
- Skin infection common in children
- Spread by direct contact
- Spread by scratching (autoinfection; virus in lesions)
- “Flesh-colored dome” lesions
- Central dimple
- Sometimes itchy
- Usually self-limited
- Resolves weeks to months




E van Herk/Wikipedia

# Herpes Viruses

Jason Ryan, MD, MPH

## DNA Viruses




All DNA viruses (+) have icosahedral capsid except Pox  
All DNA viruses replicate in nucleus except Pox

## Herpes Viruses

- All enveloped, double stranded and linear
- All have icosahedral core
- Large viruses (only Pox is bigger)
- Replicate in nucleus
- Envelope from nucleus of cell (no cell membrane)
- Often cause latent infection
  - Acute disease followed by asymptomatic period
  - Virus may reactivate later

## Herpes Viruses

- Four herpes viruses lead to giant cell formation
  - HSV1, HSV2, VZV, CMV
- Can be seen in Tzanck smear (HSV test)



CDC/Public Domain

## Herpes Viruses

- Many clinically important infections
  - Oral/genital herpes
  - Mono (EBV)
  - Chickenpox/Shingles (VZV)
  - Roseola
  - Kaposi Sarcoma
  - CMV infections

## HSV1, HSV2, VZV

- HSV-1: Oral herpes, other infections
- HSV-2: Genital herpes, neonatal herpes, meningitis
- VZV: Chicken pox, Shingles
- Many similarities
  - Primary infection phase
  - Lay dormant in nerve ganglia
  - Can reactivate
  - Treatment: acyclovir, valacyclovir, famciclovir

## HSV1

- Favors oral mucosa
- Transmitted in saliva
- Many primary infections asymptomatic
- Common initial infection: Gingivostomatitis
  - Severe sore throat
  - Painful vesicles on throat, pharynx
- Once infected, virus lives in latent state
  - Nerve cell bodies in ganglion neurons
  - Often trigeminal nerve ganglia
- Reactivation: Herpes labialis

## HSV1

Gingivostomatitis



Klaus D. Peter, Gummersbach, Germany/Wikipedia

Herpes Labialis



CDC/Public Domain

## Cold Sores

- Oral herpes often called "coldsore"
- Occur at vermillion border (edge of lips)
- Don't confuse with aphthous ulcers (canker sores)
  - Inner surfaces of lips, buccal mucosa
  - Not preceded by vesicles
  - Not caused by infection
  - Immune related



CDC/Public Domain

## HSV1

### Other Infections

- Herpetic whitlow
  - Inoculation of virus into skin break in finger
  - Painful lesion on fingertip
- Keratoconjunctivitis
  - Infection of cornea/conjunctiva
  - Pain, redness, discharge
- Encephalitis (temporal lobes)



James Heilman, MD/Wikipedia

## HSV2

- Favors genital mucosa
- Sexually transmitted
- Initial infection can be asymptomatic
- Classic symptoms: painful ulcers
- Virus enters latent phase in lumbar-sacral ganglia
- Recurrent eruptions of vesicles/ulcers
  - For frequent recurrences, suppressive Rx (acyclovir)
- Also can cause meningitis

## Congenital Herpes

### HSV2

- Newborn infection from infected mothers
- Serious infection more common when mother has primary infection
  - More virus replication
  - Fewer maternal antibodies
- Vesicular lesions on skin, eyes, mouth
- Can progress to CNS disease/encephalitis
- Seizures, poor feeding
- No congenital defects

## Varicella Zoster

VZV

- Spread through air from infected persons
- Primary infection: Chicken pox
  - Highly contagious
  - Fever, sore throat
  - Diffuse (face, trunk, limbs) vesicular rash – very itchy
  - Classic progression: macules → papules → vesicles
  - Different stages in different parts of body
  - Eventually lesions crust, fall off
- Rare complications (often adults):
  - Encephalitis
  - Pneumonia

## Varicella Zoster

VZV

- Reactivation of VZV: Herpes Zoster
- Virus will lay dormant in dorsal root ganglia
- Reactivated lesions classically follow dermatome
- Do not cross midline



Image courtesy of Fisle/Wikipedia

## Varicella Zoster

VZV

- Pre-eruptive phase (1-10 days)
  - Sensory phenomena along dermatomes
  - Pain; less commonly itching or paresthesias
  - Other (rare) symptoms: malaise, myalgia, HA, photophobia
- Eruptive phase
  - Rash with pain
  - Most commonly a thoracic dermatome
  - Lymphadenopathy may be present

## Post-Herpetic Neuralgia

- Can occur following resolution of zoster infection
- Constant or intermittent “stabbing” pain
- May last for months

## Varicella Zoster

VZV

- Age is most important risk factor
  - Rare <50 years old
- Immune compromise
  - Transplant patients
  - Immunosuppressive drugs
- Special risk group: Inflammatory bowel disease

## Varicella Zoster

VZV

- Rare complications:
  - Ophthalmic zoster (blindness)
  - Encephalitis
- Treatment:
  - Often supportive care only
  - Rarely steroids and acyclovir drugs

## Diagnosis

HSV1, HSV2, VZV

- Modern tests of choice:
  - PCR (especially CSF for encephalitis)
  - Viral culture (1 to 3 days)
  - Serology (primary infection)
- Tzanck Smear
  - Used to diagnose HSV1, HSV2, or VZV
  - Microscopic exam of scraped ulcer
  - Stained with Giemsa or Wright stain
  - Positive if multinucleated giant cells seen
- Biopsy of infected tissue
  - Can see intranuclear inclusions
  - "Cowdry A inclusions"

## EBV

- Causes mononucleosis ("mono")
- Spread by direct contact, saliva ("kissing disease")
- Virus infects and transforms B cells

## EBV

- Envelope gp350/220 binds B-cell receptor CD21
  - Receptor for C3d fragment of complement
  - Also called C3d receptor, EBV receptor, CR2


C3 → C3b → C3d

## Infectious Mononucleosis

- Classic presentation
  - Age 15-20 (college student)
  - Fever
  - Hepatosplenomegaly (splenic rupture)
  - Pharyngitis
  - Posterior cervical lymphadenopathy
  - Atypical lymphocytes (T cells)
- No specific treatment
- Most symptoms resolve weeks
- Fatigue may last months

## Atypical Lymphocyte

- Hallmark of infectious mononucleosis
- Majority are CD8+ T cells



Bobigalindo/Wikipedia

## Infectious Mononucleosis

Diagnosis

- Suggested by symptoms, lymphocytosis, atypical lymphocytes
- Heterophile antibodies ("Mono spot")
  - Heterophile antibodies agglutinate sheep or horse RBCs
  - Lab kits used
  - Sample of patient's blood (often finger stick)
  - Color change if heterophile antibodies present in plasma
  - Quick, highly specific
  - False negatives possible
- EBV-specific antibodies
  - Done when mono spot negative

## EBV Extras

- Amoxicillin rash
  - Amoxicillin given to mono patient for sore throat
  - Diffuse maculopapular rash
  - Mechanism not understood
- VDRL false positive
  - Common cause false positive VRDL
  - Don't confuse with syphilis
- After primary infection can reactivate later
  - Reactivation common in new HIV/AIDS patients

## EBV

### Other diseases

- Infection associated with many cancers
- Lymphomas
  - Burkitt, Hodgkin lymphoma, T-cell
- Nasopharyngeal carcinoma (especially China)
- Tumors in HIV patients
  - Non-Hodgkin lymphoma, Burkitt, CNS lymphoma
- Oral hairy leukoplakia
  - White plaques on tongue, cannot be scraped off
  - Classic finding in HIV patients with low CD4 count

## Cytomegalovirus

### CMV

- Ubiquitous virus
- Spread in multiple ways:
  - Sexually transmission
  - Direct contact (family, day care)
  - Blood or tissue exposure
  - Perinatal (in utero, during birth)
- Infected cells (biopsy): Owl's Eye nuclei
  - Large, dark inclusions from CMV infection
  - Perinuclear halo
- Can become latent in monocytes, marrow cells
- Treatments: Ganciclovir, Foscarnet, Cidofovir



Wikipedia/Public Domain  
Dr. Edwin Ewing, Jr., CDC

## Cytomegalovirus

### Infections

- Mostly affects immunocompromised
- Exception: CMV Mononucleosis
  - Similar to EBV infection
  - Monospot will be negative
  - Less lymphadenopathy, splenomegaly

## Cytomegalovirus

### Immunocompromised infections

- HIV, Transplant patients
- Pneumonia
  - Common after lung transplant
- Retinitis
  - Retinal edema/necrosis
  - Floaters, ↓vision
  - HIV: Low CD4 (50-100)



Wikipedia/Public Domain

## Congenital CMV

- TORCH Infection
- Most infected newborns are asymptomatic
  - Some without symptoms will develop progressive hearing loss
- Potential findings
  - Small for gestational age, microcephaly
  - Hepatosplenomegaly
  - Rashes: "Blueberry muffin syndrome"
  - Seizures
  - Sensorineural hearing loss
- Defects more common if fetus infected 1<sup>st</sup> trimester
- Treatment: ganciclovir or valganciclovir

## Newborn Deafness

### CMV

- Blueberry muffin baby
- Seizures
- Hepatosplenomegaly

### Rubella

- Blueberry muffin baby
- Cataracts
- Congenital heart disease

## HHV-6

- Causes roseola infantum (sixth disease)
- Most often due to HHV-6 but can also be caused by HHV-7 and some other viruses
- Occurs sporadically, often no exposure

## HHV-6

- Starts with febrile phase
  - High fever for several days
  - Irritable baby
  - Lymphadenopathy
  - Often confused with meningitis
- Rash
  - Fever breaks
  - Maculopapular rash
  - Starts neck and trunk
  - Spreads to face and limbs



Wikipedia/Public Domain

## HHV-8

- Causes Kaposi's sarcoma (usually HIV patients)
- Transmitted unclear
- Infects/transforms endothelial cells
- Inactivates tumor suppressor genes
- Purplish plaques/nodules on skin
  - Sometimes mouth, GI tract, lungs




M. Sand, D. Sand, C. Thrandorf, V. Paech, P. Altmeyer, F. G. Bechara

# RNA Viruses

Jason Ryan, MD, MPH

## RNA Viruses



## Caliciviruses

- Non-enveloped, (+) ssRNA, linear, icosahedral
- Norovirus genus (Norwalkvirus)
- Viral gastroenteritis
  - Recall: Most gastroenteritis is VIRAL
  - 2-3 days of watery diarrhea, nausea, vomiting
  - Not inflammatory: non-bloody, no mucous, no fecal leukocytes

## Caliciviruses

- Fecal-oral transmission
  - Often involves contaminated sea food
  - Low infectious dose, shed in stool for weeks after infection
- Commonly causes outbreaks
  - Schools (children), cruise ships, hospitals/nursing homes
- Usually diagnosed clinically, no specific treatment

## Hepeviruses

- Non-enveloped, (+) ssRNA, linear, icosahedral
- Hepatitis E virus

## Reoviruses

- Non-enveloped, dsRNA, icosahedral
- Segmented, linear viruses
- Contain RNA-dependent RNA polymerase
  - Required to make mRNA from dsRNA
- Coltivirus: Colorado tick fever
  - Transmitted by wood tick bite (*Dermacentor andersoni*)
  - Lives in rodents (squirrels, chipmunks) Rocky Mountains
  - Fever, chills, myalgias, headache
  - Self-limited
- Rotavirus

## Rotavirus

- Causes gastroenteritis in children
  - Fecal-oral transmission
  - Infects mucosal cells
  - Excess secretion of fluids, electrolytes
- Watery diarrhea
  - No blood, mucous, few/no fecal leukocytes
- Diagnosis: virus in stool
- Vaccine available
  - Live, attenuated virus (oral)
  - Given to children prior to 6 months of age

## Picornaviruses

Poliovirus, Echovirus, Rhinovirus, Coxsackievirus, Hepatitis A

- Non-enveloped, (+) ssRNA, linear, icosahedral
- Synthesize a large polypeptide
- Cleavage → viral proteins
- All transmitted fecal-oral
  - Enteroviruses
- Exception: Rhinovirus (common cold)
  - Cannot survive in stomach (acid-labile)
  - Transmitted directly via respiratory droplets

## Poliovirus

- Polio (poliomyelitis)
  - Febrile illness followed by weakness/paralysis
- Inactivated poliovirus vaccine (IPV; Salk)
  - Cannot cause vaccine-associated polio
  - Only vaccine used in US
  - Preferred vaccine in developed countries
  - **Systemic** antibody response
- Live attenuated oral polio vaccine (OPV; Sabin)
  - Some advantages in developing world
  - Cheap, easy to administer (oral)
  - Fecal-oral transmission to some unimmunized contacts
  - Triggers **local immunity** in the GI mucosa

## Picornaviruses

- Echovirus
  - Aseptic (viral) meningitis
  - 90% viral meningitis: coxsackievirus, echovirus
- Hepatitis A
- Rhinovirus
  - Viral upper respiratory illness (URI)
  - Most common virus associated with "cold" symptoms

## Coxsackievirus

- Group A & B
- Aseptic meningitis (Group A & B)
- Hand, foot, and mouth syndrome (Group A)
  - Childhood illness
  - Sore throat, oral vesicles (buccal mucosa and tongue)
  - Rash: small lesions on **hands, feet**, buttocks
- Herpangina (Group A)
  - High fever, painful mouth blisters
  - Classically in children during summer
- Myocarditis, pericarditis (Group B)

## Coxsackievirus



CDC/Public Domain

Wikipedia/Public Domain

## Coronavirus



CDC/Public Domain

- Enveloped, (+) ssRNA, linear, helical
- Upper respiratory infection ("cold")
- Severe acute respiratory syndrome (SARS)
- Worldwide outbreak in 2003
- Up to 1 week prodrome
  - Fever, malaise, headache, myalgias
- Cough, dyspnea
- Sometimes progressing to respiratory failure

## Retroviruses

- Enveloped, ssRNA, linear
- Uses reverse transcriptase to convert RNA → DNA
- DNA replicates in nucleus
- Most important example is HIV
- Other example is human T-lymphotropic virus (HTLV)

## Human T-lymphotropic virus

HTLV

- Two identical strands of (+) RNA
- Enters CD4 T-cells
- RNA genome reverse transcribed to DNA
- DNA product integrated into host cell genome
- T-cell proliferation and transformation
- Results: T-cell leukemia-lymphoma (NHL variant)

## Human T-lymphotropic virus

HTLV

- Endemic outside US (Caribbean, Africa)
- Many in US infected by IV drug use
- Infects millions, few develop leukemia
- Uncommon in US

## Flaviviruses


- Enveloped, (+) ssRNA, linear, icosahedral
- Hepatitis C
- Four mosquito illnesses
  - Yellow fever
  - Dengue fever
  - St. Louis encephalitis
  - West Nile virus

## Yellow Fever

- Occurs in Africa, South America
- Virus can live in monkeys
- Arbovirus: transmitted by mosquito bite (Aedes)
- Infects liver (yellow for jaundice)
  - High fever, headache, jaundice, high bilirubin level
- Increased AST/ALT
  - Special feature: AST >> ALT
- Can cause hemorrhage ("black vomit")
  - Coffee-ground vomit, oozing from gums
- No specific treatment
- Vaccine available

## Yellow Fever

- Councilman bodies in liver



Nephron/Wikipedia

## Dengue Fever

- Occurs in Asia, South America
- Transmitted by mosquito bite
- "Break bone fever"
- Fever with headache and retro-orbital pain
- SEVERE muscle and joint pains
- Maculopapular rash
- Can rarely progress to hemorrhagic shock
- No specific therapy

## St. Louis and West Nile

- Both transmitted by mosquito bites (arboviruses)
- Birds carry the virus
- Both cause encephalitis
  - Fever, altered mental status
  - Sometimes meningitis symptoms (stiff neck, photophobia)
- Most people asymptomatic
- St. Louis:
  - Widely distributed in the Americas
  - Few cases per year, sometimes outbreaks in US
- West Nile:
  - Widely distributed across the globe
  - Outbreaks have occurred in US

## Togaviruses

- Enveloped, (+) ssRNA, linear, icosahedral
- Rubella
- Eastern equine encephalitis
- Western equine encephalitis

## Rubella

German measles; 3-day measles

- Childhood exanthem (rash)
- Acquired by inhalation of respiratory droplets
- Mild fever, lymphadenopathy
- Maculopapular rash (1-5 days after fever)
- Characteristic lymphadenopathy
  - Posterior cervical
  - Posterior auricular
- No specific treatment
- Vaccine: Live attenuated virus (MMR)

## Congenital Rubella Syndrome

- TORCHES infection
- Mother acquires infection via respiratory droplets
  - Rash, fever, lymphadenopathy
- Classic triad in fetus:
  - Deafness
  - Cataracts
  - Cardiac disease

## Congenital Rubella Syndrome

- Congenital heart disease
  - Patent ductus arteriosus (PDA)
  - Pulmonary artery stenosis
  - Many, many others
- Also petechiae/purpura ("blueberry muffin baby")
- Babies excrete virus for months
- Public health hazard
- Diagnosis:
  - IgM antibodies (recent infection)
  - Amniocentesis (virus in amniotic fluid)

## Newborn Deafness

- | CMV                     | Rubella                    |
|-------------------------|----------------------------|
| • Blueberry muffin baby | • Blueberry muffin baby    |
| • Seizures              | • Cataracts                |
| • Hepatosplenomegaly    | • Congenital heart disease |

## Equine Encephalitis

- Eastern and Western Equine Encephalitis
- Both can infect humans and horses (equine)
- Found in North America
- Virus resides in birds
- Transmitted by mosquitos
- Most infections asymptomatic
- Can cause encephalitis
  - Fever, mental status changes

## Bunyaviruses

- Enveloped, (-) ssRNA, helical
- Segmented (BOAR), circular genome (BAD)
- Result in rare infections
- California encephalitis
  - Mosquito-borne virus, causes encephalitis
  - Reservoir is rodents
- Hemorrhagic fever
  - Rift Valley fever
  - Crimean-Congo hemorrhagic fever
  - Hantavirus infection

## Encephalitis Viruses

Mosquito-borne Arboviruses

- Flaviviruses (birds)
  - St. Louis
  - West Nile
- Togaviruses (birds)
  - Eastern Equine
  - Western Equine
- Bunyaviruses (rodents)
  - California

## Hemorrhagic Fever

- Viral infections by enveloped RNA viruses
- Live in animals, usually birds
- Most transmitted in one of two way:
  - Mosquito or tick bites
  - Contact with infected animals
- Initial symptoms non-specific
  - Fever, headache, malaise
  - Often GI symptoms: vomiting, diarrhea
- Hemorrhage may occur
  - Petechiae, large hematomas, frank bleeding
- Can progress to respiratory failure, shock, death

## Bunyaviruses

### Hemorrhagic Fever

- Rift Valley fever
  - Mosquito-borne virus, East Africa
  - Transmitted by mosquito bite or contact infected animals
- Crimean-Congo hemorrhagic fever
  - Tick-borne virus, East Africa
  - Transmitted by ticks or contact infected livestock
- Hantavirus infection
  - Lives in rodents (mice)
  - Transmitted by rodent contact
  - Virus shed in rodent urine, feces, saliva
  - Often progresses: renal failure or respiratory failure

## Filoviruses

- Enveloped, (-) ssRNA viruses, linear, helical
- Transmitted through contact with body fluid from infected person
- Ebola & Marburg
- Both cause hemorrhagic fever
- Both highly fatal

## Arenaviruses

- Enveloped, (-) ssRNA, helical
- Segmented (BOAR), circular genome (BAD)
- Lassa fever
  - Hemorrhagic fever
  - Spread by urine from rats
  - Also through close contact with infected people
- Lymphocytic choriomeningitis virus (LCMV)
  - Rare cause of viral meningitis
  - Rats and mice shed virus in saliva, urine, feces
  - Children in poor conditions at higher risk exposure

## Hemorrhagic Fever Viruses

- Bunyaviruses
  - Rift Valley Fever virus (mosquito)
  - Crimean-Congo hemorrhagic fever virus (tick)
  - Hanta virus (rodents)
- Filoviruses
  - Ebola/Marburg
- Arenaviruses
  - Lassa fever (rats)
- Sometimes flaviviruses
  - Yellow fever, dengue can progress

## Rhabdoviruses


- Enveloped, (-) ssRNA, linear, helical
- Several rhabdo species cause rabies

## Rabies

- Infection by bite of rabid animal
- Found in bats, raccoons, skunks, coyotes, wolves
- Classic transmission from dog bite
- Incubation period: 1 to 3 months after bite
- Prodrome: Fever, malaise, nausea, vomiting
- Rabies infection
  - Encephalitis
  - Painful pharyngeal spasms
  - Classically fear of water, agitation, salivation
- Progresses to paralysis, coma
- Virtually always fatal

## Rhabdoviruses

- Special features of rabies viruses
  - Bullet shaped envelope
  - Forms "Negri bodies" in neurons/Purkinje cells
  - Viruses bind Ach receptors on peripheral nerves
  - Migrate to CNS



Wikipedia/Public Domain

## Rabies

### Management of Possible Infection

- Suspect rabies after:
  - Wild animal bite
  - Exposure to bats
- Important to clean bite wounds
- Vaccination: Inactivated virus vaccine
- Rabies immune globulin
  - Provides antibodies until protective antibodies generated from vaccination

## Orthomyxoviruses

- Enveloped, (-) ssRNA, linear, helical
- Segmented genome (BOAR)
- Most important virus in influenza

## Influenza Virus

- Causes acute respiratory illness (the flu)
- Occurs in winter months
- Transmitted by respiratory secretions
  - Infected person's cough, sneeze
- Fever, headache, myalgia, and malaise
- Cough, sore throat, runny nose
- Self-limited; improvement in days to weeks
- Rare complications
  - Pneumonia (viral or secondary bacterial)
  - S. pneumoniae most common; S. aureus 2<sup>nd</sup> most common

## Influenza Virus

- Several subtypes (A, B, C)
- Influenza A most common
- Two key envelope glycoproteins
- Hemagglutinin
  - Binds to cells; assists in entry into cells
- Neuraminidase
  - Remove sialic acid from glycoproteins
  - Required for virus progeny to exit cell
- Virus replicates in nucleus

## Influenza Virus

- Antigenic drifts
  - Minor changes in hemagglutinin and neuraminidase
  - Due to random mutation
  - Vary from year to year
  - Some previously infected not immune (epidemics)
- Antigenic shift
  - Segmented genome allows for high rates of reassortment when two viruses infect same cell
  - Often occur when animal/human virus infect same cell
  - Cause pandemics (US in 1918)
  - Can occur in other segmented viruses
  - BOAR: Bunyavirus, arenavirus, orthomyxovirus, reovirus

## Influenza Vaccine

- Viruses grown in eggs
- WHO recommends strains for vaccine
  - Global surveillance of viruses at end of prior flu season
- Killed vaccine virus
  - Available each fall (prevents winter flu)
- Nasal spray
  - Live, temperature sensitive mutant
  - Replicates in nose, not lungs
  - Often used for children (cannot use >age 50)

## Paramyxoviruses

- Enveloped, (-) ssRNA, helical
- All cause disease in children
- All contain F protein
  - Surface F (fusion) protein
  - Causes respiratory epithelial cell fusion
  - Palivizumab: monoclonal antibody against F protein
  - Used to treat RSV

## Paramyxoviruses

- Parainfluenza
  - Croup
  - URI in children with "barking" cough (sounds like a seal)
  - Can cause respiratory distress (treat with steroids)
- RSV
  - Viral respiratory infection in infants
  - Treatment: Ribavirin, Palivizumab
- Measles
- Mumps

## Measles

Rubeola

- Cough, Coryza, Conjunctivitis
- Classic maculopapular rash
  - Starts at head → spreads to feet
- Koplik spots
  - Small, white lesions in mouth
- Rare complications
  - Measles encephalitis
  - Subacute sclerosing panencephalitis (SSPE) – YEARS after
    - Personality changes, odd behavior, dementia
  - Giant cell pneumonia
    - Immunocompromised
    - Multinucleated giant cells in lung tissue

## Measles



Wikipedia/Public Domain



Wikipedia/Public Domain

## Measles

- 2014 outbreaks in US among unvaccinated children
- Can spread to vaccinated children
  - Vaccine 95% effective
- Test of choice:
  - Measles IgM
  - Not positive first few days of infection
- Possible therapies:
  - Vitamin A
  - Ribavirin

## Mumps

- Prodrome of fever, malaise, headache, myalgias
- Parotitis
  - Inflammation of parotid glands (facial swelling)
- Orchitis
  - Testicular pain
  - Scrotal swelling
  - Can result in sterility
- Meningitis (aseptic)



Wikipedia/Public Domain

## MMR Vaccine

- Measles, Mump, Rubella
- Live, attenuated vaccines
- Usually given after 1-year
- Prior to 1-year, maternal antibodies will kill vaccine
- Live, attenuated vaccine required:
  - Paramyxoviruses (measles, mumps) form "syncytia"
  - Move from cell to cell directly
  - No exposure to plasma (antibodies)
  - Need vigorous cell-mediated response to infection

## Childhood Red Rashes

Virus	Features
Rubella (German Measles)	Head → Feet Postauricular LAD
Measles (Rubeola)	Head → Feet Cough, Coryza, Conjunctivitis Koplik spots
HHV-6 (Roseola)	Fever that breaks Neck/Trunk - Face/limbs Infants/Seizures
Parvovirus B19	Slapped cheek
Strep Pyogenes (Scarlet Fever)	Sore throat Sandpaper-like red rash Head/neck → Trunk

# Viral Hepatitis

Jason Ryan, MD, MPH

## Viral Hepatitis

- Hepatitis viruses (A, B, C, D, E)
- All cause liver inflammation
- Some cause chronic infection
- Can lead to cirrhosis or hepatocellular carcinoma

## Viral Hepatitis

### Acute Symptoms

- Many acute infections asymptomatic
- Fever, malaise, nausea, vomiting, anorexia
- RUQ pain
- Jaundice (yellow skin from bilirubin)
- Itching (bile salts in skin)
- Dark urine (bilirubin)
- Clay-colored stools (lack of bilirubin excretion)

## Viral Hepatitis

### Acute Symptoms

Conjugated  
Bilirubin  $\xrightarrow{\text{Bacteria}}$  urobilinogen  $\rightarrow$  stercobilin  $\rightarrow$  

Nemo/CCO Public Domain

## Blood Tests

- Increased AST/ALT
  - ALT usually > AST
  - Contrast with alcoholic hepatitis (AST>ALT)
- Increased bilirubin (direct)
  - Liver can conjugate bilirubin in setting hepatitis
  - Cannot transport into bile
- False positive VDRL
  - Viral hepatitis is common cause of false positive VDRL
  - Don't confuse with syphilis

## Hepatitis A

- Picornavirus (PERCH)
- Non-enveloped, (+) ssRNA, linear, icosahedral
- Synthesize a large polypeptide
- Cleavage  $\rightarrow$  viral proteins
- All transmitted fecal-oral

## Hepatitis A

- Transmitted through:
  - Personal contact
  - Drinking contaminated water
  - Consumption of raw sea food
- Common in underdeveloped countries
  - Poor hygiene and sanitation
- Classic case: traveler to Mexico, Central/South America
- Incubation period ~30 days

## Hepatitis A

- Diagnosis:
  - Acute disease: Anti-HAV IgM antibodies plus symptoms
  - Prior disease: Anti-HAV IgG antibodies
- Self-limited; no specific therapy
- Acute disease only – no chronic infection
- Often asymptomatic
  - Antibody tests done later may show anti-HAV IgG
- Inactivated virus vaccine available (IM)
- Part of US routine childhood vaccination schedule

## Hepatitis E


- Hepevirus
- Non-enveloped, (+) ssRNA, linear, icosahedral
- Outbreaks worldwide in resource-limited areas
- Infection from fecal contamination of water
- Self-limited acute infection - no chronic infection
- Diagnosis:
  - HEV genome in in serum or feces (PCR)
  - IgM antibodies to HEV
- Pregnancy
  - Hepatic failure more frequent during pregnancy
  - High mortality rate (15 to 25 %)

## Hepatitis B

- Hepadnavirus family (DNA virus)
- Enveloped, circular, icosahedral capsid
- #1: Partially double stranded DNA virus
  - Genome enters hepatocytes → nucleus
  - DNA becomes fully double stranded
  - mRNA synthesized → cytoplasm
- #2: Reverse transcriptase synthesized
  - Viral mRNA → viral DNA
  - Packaged in capsid
- #3: Envelope from endoplasmic reticulum


## Hepadnavirus

Hepatitis B virus



## Hepatitis B

Dane Particle



Public Domain/Wikipedia

## Hepatitis B

### Transmission

- Sexual contact
- IV (drug use, transfusion, needle stick)
- Maternal-fetal
  - Especially if mother gets acute disease 3<sup>rd</sup> trimester
  - Babies usually have minimal symptoms
  - Lots of viral replication in baby (immature immune system)
  - Babies at HIGH risk of progression to chronic disease
- Incubation of 1 to 4 months
- Acute infection
  - 70% have subclinical or mild hepatitis (anicteric)
  - 30% icteric hepatitis

## Chronic Hepatitis B

- Progression to acute → chronic depends on age
  - 90% peri-natal
  - ~50% children
  - <5% adults
- Many chronic infections asymptomatic (carriers)
- Risk of progression to:
  - Cirrhosis
  - Liver failure
  - Hepatocellular carcinoma (viral DNA integrates into host)
  - Reactivation (acute hepatitis)

## Immune Reconstitution

- If HIV is treated without treating Hep B it can cause severe liver damage
- HepB testing usually done prior to HIV therapy

## Extrahepatic Manifestations

- Polyarteritis nodosa
  - Fevers, fatigue, arthralgias
  - Abdominal pain, melena
  - Neuropathy
  - Rash
- Glomerular disease
  - Most common is membranous nephropathy
  - Presents as proteinuria, nephrotic syndrome

## Hepatitis B

### Diagnosis

- Antigenes
  - Hepatitis B surface antigen (HBsAg)
  - Hepatitis B e antigen (HBeAg)
- Antibodies
  - Anti-hepatitis B surface antigen (Anti-HBsAg)
  - Anti-hepatitis B e antigen (Anti-HBeAg)
  - Anti-hepatitis B core antigen (Anti-HBcAg)
- Antigens rise in acute disease, fall as infection resolves
- Antibodies rise as acute infection resolves

## Hepatitis B Surface Antigen

### HBsAg

- Hallmark of infection
- Glycoprotein that forms spheres and tubules (EM)
- From surface of envelope
- Detectable weeks after exposure, prior to symptoms

## Hepatitis B Surface Antigen

HBsAg

- Recovery from acute hepatitis:
  - HBsAg becomes undetectable after four to six months
- Chronic infection
  - Persistence of HBsAg for more than six months
- Prior infection or vaccination:
  - Presence of anti-HBsAg antibodies without HBsAg
- Generally, when anti-HBsAg levels rise, HBsAg levels fall and infection clears

## Hepatitis B Vaccine

- Contains recombinant HBsAg
- Vaccinated individuals will be (+) anti-HBsAg
- All other antibodies (HBc, HBe) should be negative

## Hepatitis B Core Antigen

HBcAg

- Intracellular antigen (comes from within hepatocytes)
- Capsid core protein
- Expressed by infected hepatocytes
- NOT detectable in serum
- Anti-HBc can be detected
- Anti-HBc IgM rises in acute infection
- Anti-HBc IgG prior exposure or chronic infection

## Window Period

- Brief period where:
  - HbsAg undetectable
  - Anti-HBsAg not yet detectable
- Can give false appearance of no infection
- SOLE marker of infection is anti-HBc (IgM)

## Hepatitis B e Antigen


HBeAg

- Viral protein secreted by infected cells
- Part of capsid core
- Indicates significant viral replication
  - Correlates well with levels of HBV DNA
  - HBsAg indicates presence of virus, not necessarily significant replication
- Elevated in patients who are **highly infectious**
- Seroconversion to Anti-HBeAg usually associated with fall in viral DNA production

## Hepatitis B DNA

- Detectable with PCR
- Major role is for determining "viral load" for treatment

## Hepatitis B Diagnosis



## Hepatitis B Diagnosis

- Acute disease
  - HBsAg (except window)
  - IgM Anti-HBc (even during window)
  - HBeAg (indicates infectivity)
  - HBV DNA
- Recovery after acute disease
  - Anti-HBs; Anti-HBe; Anti-HBc (IgG)
  - Undetectable HBsAg
  - Absence of HBV DNA

## Diagnosis of Hepatitis B

- Prior vaccination
  - Anti-HBsAg only
  - Not anti-HBc or anti-HBe
- Chronic infection
  - HBsAg positive
  - If HbeAg positive = high infectivity
  - Anti-HBc positive
  - Viral DNA may be high, low depending on viral load
- If HBsAg is positive = patient is infected
- If HBsAg is negative = patient is not infected

## Treatment Chronic Hepatitis B

- Acute hepatitis B
  - Usually treated with supportive care
  - Immunocompetent adults <5% chance chronic disease
- Multiple treatments for chronic disease
  - Interferon
  - Lamivudine (NRTI)
  - Other antiviral drugs

## Interferons

- Type I: Alpha, beta (most human cells)
- Type II: Gamma (T cells, NK cells)
  - Increase MHC expression, activates NK cells
- Activate interferon stimulated genes (ISGs)
  - Many, many cellular effects

## Interferons

- Trigger cell production of RNAase L
  - Degrades viral and cellular RNA
- Production of protein kinase R
  - Inhibits translation of proteins
- Active only in presence dsRNA

## Interferons

- Administered as drugs
- Antiviral: Hepatitis B, C (alpha)
- Leukemia, Lymphoma, Melanoma (alpha, beta)
- Multiple sclerosis (beta)
- Gamma:
  - Rare immune diseases
  - Chronic granulomatous disease (CGD)
  - IL-12 receptor deficiency

## Hepatitis C

- Flavivirus (mostly mosquito illnesses)
- Enveloped, (+) ssRNA, linear, icosahedral
- High degree of antigenic variation
- Envelope glycoproteins
  - Contain a "hypervariable region"
  - High mutation rate in genome
  - Lack of proofreading by viral RNA polymerase
  - Result: prone to frequent mutations
- Difficult for immune system to eradicate effectively
- High rate of chronic disease

## Hepatitis C

- Mostly acquired through IVDA or transfusion
  - Transfusion illness now rare due to screening
- Rare cases from needle sticks, sexual contact
- Acute illness
  - Usually asymptomatic
  - Usually leads to chronic disease
- Chronic infection
  - Usually asymptomatic or mild, nonspecific symptoms
  - Often incidental discovery of abnormal LFTs
  - Screening done for high risk patients (IVDA)

## Hepatitis C

### Diagnosis

- HCV RNA by PCR
  - Elevated soon after exposure
- Anti-HCV
  - Elevated by 12 weeks after exposure
- Both elevated in chronic disease (common)

## Hepatitis C

- Chronic infection associated with:
  - Cirrhosis
  - Liver failure (common indication for transplant)
  - Hepatocellular carcinoma
- Treatment options:
  - Interferon
  - Ribavirin
  - Others

## Hepatitis D

### Delta Agent

- Small enveloped (-) RNA virus, circular genome
- "Defective virus"
- Lacks genes for envelope proteins
- Uses HbsAg for envelope protein
- Genome encodes one protein: delta antigen (HDAg)
- Virus particles carry HDAg

## Hepatitis D

### Delta Agent

- Pathogenesis:
  - Invades hepatocytes
  - Travels to nucleus to replicate
  - Uses HBV to provide envelope
  - Virus particle coated with HBsAg
  - Uses host cell RNA polymerase to replicate genome

## Hepatitis D

### Delta Agent

- Transmission:
  - Co-infected with HBV
  - HDV infection in setting of chronic HBV carrier state
  - Superinfection often leads to flare of hepatitis
- Diagnosis:
  - Serum HDAg
  - HDV RNA
  - Anti-HDV antibodies
- Hep B vaccine protects against Hep D

# HIV

Jason Ryan, MD, MPH

# HIV

Human Immunodeficiency Virus

- RNA retrovirus
- Uses reverse transcriptase: RNA → DNA
- Infects CD4+ T-helper cells
- Acquired immunodeficiency syndrome (AIDS)
- Susceptibility to unique opportunistic infections
- Natural host is humans


# HIV

- Diploid: Carries two copies of (+) stranded RNA
- RNA tightly bound to key enzymes
  - Reverse transcriptase: Makes DNA from RNA
  - Aspartate protease: Cleavage of proteins
  - Integrase: Integrate DNA into host cell DNA
- Cone-shaped (conical) capsid
  - Multiple copies of p24 protein
  - Non-variable protein
  - Antibodies produced but do not neutralize virus

# HIV

- Enveloped
  - Phospholipid from membrane of human cell
- Envelope contains Env protein
- Cap of env protein contains glycoproteins
  - gp120: Attachment to T-cells
  - gp41: Fusion and entry into T-cells
  - Both formed as single protein (gp160) and cleaved
- Gene for gp120 mutates rapidly (antigenic variation)
  - "V3 loop" portion is highly immunogenic
  - Varies significantly
  - Antibody neutralization difficult

# HIV




Wikipedia/Public Domain

# HIV Genome

- Three "main" genes encode major parts of virus
- Gag: nucleocapsid
- Pol: polymerase
- Env: envelope proteins

## HIV Genes and Products



## Other HIV Genes


### Regulatory and Auxiliary Genes

- Tat, Rev
  - Required for viral replication
  - Tat: Activates transcription of genes
  - Rev: Transports mRNA from nucleus to cytoplasm
- Nef, vif, vpr, vpu, vpx
  - Not required for replication (auxiliary)
  - Nef: ICD4 proteins and MHC I on T-cell surfaces

## HIV Tropism

- CCR5 and CXCR4
  - Major lymphocyte receptors used by HIV to enter cells
- CCR5-tropic viruses
  - Replicate in monocytes/ macrophages (M-tropic)
  - Can also infect dendritic cells (trafficking to lymph nodes)
  - Occur early: Sexually transmitted to macrophages
- CXCR4-tropic viruses
  - Replicate more efficiently in T-cells (T-tropic)
  - Occur later after infection has developed
- Mutations in CCR5 gene associated with decreased susceptibility to HIV infection

## Pathophysiology



Wikipedia/Public Domain

## HIV-1 and HIV-2

- Two types of HIV (two viruses) cause infection
- HIV-1: Causes majority of infections worldwide
- HIV-2: Important cause of infection West Africa
- Both sexually transmitted
- Both can cause AIDS
- Some differences in progression, severity
- One drug class (NNRTIs) not effective HIV-2

## HIV Transmissions

- Sexually transmitted
- Exposure to contaminated blood
- Perinatal transmission


## Markers of Infection

- CD4 T-cell count
  - Determined by flow cytometry
  - Normal  $\sim 1000$  cells/mm<sup>3</sup>
  - AIDS  $< 200$
  - Used to initiate prophylaxis against opportunistic infections
- Viral load
  - Quantification of HIV RNA
  - Used to monitor effect of drug therapy

## HIV Symptoms

- Initial infection asymptomatic 10-60%
- Acute HIV syndrome
  - 2-4 weeks after exposure
  - Fever, myalgias, sore throat, cervical adenopathy
  - Sometimes maculopapular rash
  - Similar to mononucleosis
- Severe immunosuppression (AIDS)
  - Average time of 8 years from exposure
  - CD4  $< 200$  cells/mm<sup>3</sup> or AIDS-defining infection
  - Symptoms from opportunistic infections

## HIV Time Course



## Rare HIV Features

- Dementia
- Pulmonary hypertension
- Cardiomyopathy

## HIV Diagnosis

- Older tests (antibody only)
  - ELISA: Screening (sensitive; many false positives)
  - Western blot: Confirmatory if positive ELISA (specific)
- Current CDC recommendations:
  - Combination antigen/antibody tests
  - "4<sup>th</sup> generation tests"
- Test for p24 antigen and HIV antibodies
- If positive, HIV1-HIV2 antibody differentiation assay

## HIV Diagnosis

- Acute HIV
  - HIV RNA testing (viral load) will be high
- Perinatal HIV
  - Maternal HIV antibodies persist for months
  - Standard test is HIV PCR testing

## Opportunistic Infections

- Pneumocystis (fungal) pneumonia
- CMV retinitis
- Cryptococcal (fungal) meningitis
- Toxoplasmosis (CNS protozoa infection)
- Cryptosporidium diarrhea (protozoa)
- Kaposi Sarcoma (HHV-8)
- Mycobacterium avium complex
- Thrush (Candida - fungus)

## Opportunistic Infections

- Prophylaxis given based on CD4 count
- CD4 < 200
  - TMP-SMX: Pneumocystis pneumonia
- CD4 < 100
  - TMP-SMX: Pneumocystis and Toxoplasmosis
  - Itraconazole: Histoplasmosis (endemic areas)
- CD4 < 50
  - Azithromycin: Mycobacterium avium complex

## HIV Treatment

- Older guidelines based on CD4 count
  - Treat patients <500cells/mm<sup>3</sup>
- Newer guidelines recommended treating all patients
- Multi-drug therapy used (often different classes)
  - Highly active antiretroviral therapy (HAART)
- Gene mutations occur over time due to drugs
- Require altering medical regimen

## IRIS

Immune Reconstitution Inflammatory Syndrome

- Treatment of HIV → flare of infectious symptoms
- Sometimes previously undiagnosed infection
- Leading agents:
  - Tuberculosis
  - Mycobacterium avium complex (MAC)
  - Cytomegalovirus (CMV)
  - Cryptococcal meningitis
  - Pneumocystis
  - HSV
  - Hepatitis B
  - HHV-8 (Kaposi Sarcoma)

# HIV Drugs

Jason Ryan, MD, MPH

## HIV Therapy


- Protease inhibitors
- Nucleoside reverse transcriptase inhibitors (NRTIs)
- Non-nucleoside rev. transcriptase inhibitors (NNRTIs)
- Other drugs

## Protease Inhibitors

Lopinavir, Ritonavir, Indinavir

- Inhibit HIV protease
  - Product of pol gene
  - Cleaves polypeptides into smaller, functional units
  - Block production: reverse transcriptase, protease, integrase, structural proteins
- Viral particles cannot “mature”
- Become noninfectious

## HIV Genes and Products




## Protease Inhibitors

Lopinavir, Ritonavir, Indinavir

- Many side effects
  - Nausea, diarrhea
  - Hyperlipidemia, Hyperglycemia (insulin resistance)
  - Fat redistribution
- Indinavir
  - Kidney stones (hydration important)
- Ritonavir
  - Inhibits cytochrome p450 system
  - Low dose (less side effects) used to “boost” other PIs
  - Primary use of this drug is for boosting
  - Ritonavir/Lopinavir = Kaletra


## NRTIs

Nucleoside reverse transcriptase inhibitors



## NRTIs

Nucleoside reverse transcriptase inhibitors





## NRTIs

Nucleoside reverse transcriptase inhibitors


- Zidovudine, Lamivudine, Tenofovir, Didanosine
- Similar to nucleotides (ACGT)
- Lack -OH group: terminates DNA chain
- Inhibit reverse transcriptase

## Zidovudine




## NRTIs

Nucleoside reverse transcriptase inhibitors



## Nucleoside vs. Nucleotide


- Nucleotide
  - Nitrogenous base
  - Sugar
  - Phosphate group
- Nucleoside
  - Base and sugar
  - No phosphate group



## NRTIs


Nucleoside reverse transcriptase inhibitors

- Tenofovir
  - Nucleotide (contains 1 phosphate)
  - Becomes tri-phosphorylated
  - Inhibits RT as mimic of adenosine
- Zidovudine, Lamivudine, others
  - Nucleosides
  - Must be tri-phosphorylated



## NRTIs

Nucleoside reverse transcriptase inhibitors



## NRTIs

Nucleoside reverse transcriptase inhibitors

- Mitochondrial toxicity
  - Adverse effect of NRTI class
  - DNA polymerase gamma inhibited (mitochondrial enzyme)
  - Loss of mitochondria
- Symptoms:
  - Peripheral neuropathy (pain, paresthesias)
  - Myopathy
  - Pancreatitis
  - Lactic acidosis

Source: Sex Transm Infect. Jun 2001; 77(3): 158-173.

## Lactic Acidosis

- Inhibition of oxidative phosphorylation
- Anaerobic metabolism
- Lactic acidosis
- Cases of severe, life-threatening lactic acidosis reported with NRTIs

## Zidovudine

- First antiretroviral medication used for HIV
- Bone marrow suppression
- Can be improved with G-CSF or EPO
- Can be given to prevent maternal-fetal infection
  - Prenatally to mothers
  - Also given to infant
  - New WHO guidelines recommend multi-drug combinations to prevent transmission

## NRTIs

Nucleoside reverse transcriptase inhibitors

Drug	Nucleotide	Comments
Zidovudine (ZDV)	Thymidine	Bone marrow ↓
Emtricitabine (FTC)	Cytidine	
Abacavir	Guanosine	Fever, rash
Didanosine (ddI)	Guanosine	Pancreatitis
Stavudine (d4T)	Thymidine	Lipodystrophy
Lamivudine (3TC)	Cytidine	Least toxic; Hep B
Tenofovir	Adenosine	GI upset

## NRTIs


Nucleoside reverse transcriptase inhibitors

- Tenofovir
  - Fanconi syndrome
  - Loss of proximal tubule function
  - Proteinuria, urinary phosphate wasting, glycosuria
  - Metabolic acidosis, hypophosphatemia, hypokalemia
  - Polyuria, muscle weakness
- [www.aidsinfo.nih.gov](http://www.aidsinfo.nih.gov)
  - Guidelines for the Use of Antiretroviral Agents in HIV-1-Infected Adults and Adolescents

## NNRTIs

Non-nucleoside reverse transcriptase inhibitors

- Nevirapine, Efavirenz, Delavirdine
- Inhibit reverse transcriptase (different site NRTIs)
- Do not require phosphorylation
- Do not suppress bone marrow (only effect RT)
- Not effective for HIV-2 (West Africa)



Wikipedia/Public Domain

## NNRTIs

Non-nucleoside reverse transcriptase inhibitors

- GI upset
- Skin rash (rarely severe SJS)
- Metabolized by P450 system
  - Nevirapine: Inducer
  - Delavirdine: Inhibitor
  - Efavirenz: Mixed

## Other HIV Drugs

- Raltegravir
  - Integrase inhibitor
  - Integrase inserts viral DNA into cellular genome
  - Loss of activity disrupts viral life cycle
- Enfuvirtide
  - Binds gp41
  - Inhibits fusion/entry HIV
- Maraviroc
  - Blocks CCR5 on macrophages

# Antivirals

Jason Ryan, MD, MPH

## Antiviral Drugs


- Influenza drugs
  - Zanamivir, oseltamivir
- Ribavirin
  - RSV, Hepatitis C
- Acyclovir
  - Herpes viruses
- CMV Drugs
  - Ganciclovir
  - Foscarnet
  - Cidofovir
- Interferon

## Zanamivir/Oseltamivir

- Used for treatment of influenza
- Inhibit neuraminidase
  - Enzyme that cleaves sialic acid from glycoproteins
  - Required step in exit from infected cells
- Efficacy only demonstrated 1<sup>st</sup> 48hrs of illness

## DNA/RNA Drugs


General Principles



Madeleine Price Ball/Wikipedia

## DNA/RNA Drugs

General Principles




## DNA/RNA Drugs

General Principles

- Many antivirals mimic nucleotides (A, C, G, T)
  - Ribavirin, Acyclovir, Ganciclovir, Cidofovir
- Drug used by DNA or RNA polymerase
- Once used, chain terminates (inhibition)
- "Inhibitors" of RNA/DNA polymerase
- Drugs often need to become tri-phosphorylated
- Viral kinase and/or cellular kinases
- Mutations of viral kinases often lead to resistance




## Ribavirin

- Inhibition of RNA polymerase
  - Triphosphorylated by cellular kinase enzymes
  - Binds RNA polymerase, prevents binding correct nucleotides
  - Result: ↓ in viral replication/production of defective virions



## Ribavirin

- Inhibits IMP dehydrogenase
  - Inosine monophosphate dehydrogenase
  - Used to synthesize guanine nucleotides
  - Inhibited by Ribavirin
  - Decreases pool of available guanine nucleotides



Wikipedia/Public Domain


cacyle/Wikipedia

## Ribavirin

- Two main modern uses
- RSV in children
- Hepatitis C
  - Often with interferon
- Key side effect: Hemolytic anemia
  - Drug accumulates in RBCs
  - Drug phosphorylation → relative ATP deficiency
  - Hemolytic anemia in ~10% patients
  - Can be severe
- Highly teratogenic

## Acyclovir

- Inhibitor of herpes virus DNA polymerase
- Mimics guanosine → terminates chain growth



Self/Wikipedia

## Acyclovir

- Phosphorylated by **herpes virus** thymidine kinase
  - Becomes acyclovir monophosphate
  - Only occurs in infected cells (targeted effect)
- Monophosphate → triphosphate by cellular enzymes
- Acts as analog to deoxyguanosine triphosphate (dGTP)
- Inhibits viral DNA polymerase
- Resistance:
  - ↓ viral thymidine kinase
  - Altered viral thymidine kinase
  - Altered viral DNA polymerase (↓ binding acyclovir triphos.)

## Acyclovir

- Famciclovir: Similar mechanism
  - Longer half-life (lower dose can be used)
- Valacyclovir: Pro-drug, converted to acyclovir
  - Greater bioavailability (lower dose can be used)
- All 3 drugs generally well tolerated
- Acyclovir: Nephrotoxicity (IV form)
  - Crystallizes in urine
  - Given with IV fluids

## Acyclovir

- Effective for HSV-1, HSV-2, and VZV
- Uses
  - Genital herpes
  - Herpes labials
  - Herpes encephalitis
  - Herpes zoster
- Sometimes given for “suppressive” therapy

## CMV Drugs

- Used to treat CMV infections
  - HIV/AIDS
  - Transplant patients
- Three key drugs
  - Ganciclovir
  - Foscarnet
  - Cidofovir
- All interfere with CMV DNA polymerase

## Ganciclovir


- Similar mechanism to acyclovir (analog to guanosine)
- Intracellular conversion by CMV viral kinase
  - Becomes ganciclovir 5'-monophosphate
- Monophosphate → triphosphate by cellular enzymes
- Acts as analog to deoxyguanosine triphosphate (dGTP)
- Incorporation terminates chain growth

## Ganciclovir

- Major toxicity:
  - Bone marrow suppression especially leukopenia
  - Inhibits bone marrow DNA polymerase
- Valganciclovir
  - Pro-drug
  - Converted to ganciclovir
  - Better bioavailability
- Ganciclovir given primarily IV (poor bioavailability)
- Oral valganciclovir preferred for oral dosing

## Foscarnet

- Pyrophosphate analog
- Binds/inhibits viral DNA polymerase
- Blocks cleavage pyrophosphate from triphosphates
- Stops DNA chain elongation




Wikipedia/Public Domain

## Foscarnet

- Uses:
  - CMV when Ganciclovir fails
  - Acyclovir-resistant HSV and VZV
- Side effects
  - Nephrotoxicity (limiting side effect)
  - Chelates calcium (hypocalcemia)
  - Induces renal wasting of magnesium (hypomagnesaemia)
  - Seizures (often related to electrolytes)

## Cidofovir

- Nucleotide analog (cytidine)
- Cellular phosphorylation
  - No viral kinase required
- Inhibits viral DNA polymerase
- Main use is CMV retinitis
- Main toxicity is renal failure
  - Co-administer with saline
  - Probenecid (blocks renal tubular secretion of drug)



Wikipedia/Public Domain

## Interferons

- Cytokines
- Glycoproteins synthesized by infected cells
- Numerous immunomodulatory effects
- Interferon  $\alpha$ 
  - Hepatitis B and C
  - Kaposi sarcoma (HHV-8)